

Passage-time moments and hybrid zones for the exclusion-voter model

Iain M. MacPhee¹

Mikhail V. Menshikov¹

Stanislav Volkov²

Andrew R. Wade^{*,2}

2nd October 2008

¹ Department of Mathematical Sciences, University of Durham, South Road, Durham DH1 3LE, UK.

² Department of Mathematics, University of Bristol, University Walk, Bristol BS8 1TW, UK.

Abstract

We study the non-equilibrium dynamics of a one-dimensional interacting particle system that is a mixture of the voter model and exclusion process. With the process started from a finite perturbation of the ground-state Heaviside configuration consisting of 1s to the left of the origin and 0s elsewhere, we study the relaxation time τ , that is, the first hitting time of the ground-state configuration (up to translation). In particular, we give conditions for τ to be finite and for certain moments of τ to be finite or infinite, and prove a result that approaches a conjecture of Belitsky *et al.* [*Bernoulli* **7** (2001) 119–144]. Ours are the first non-existence of moments results for τ for the mixture model. Moreover, we give almost-sure asymptotic results on the long-term evolution of the size of the hybrid (disordered) region. Most of our results pertain to the discrete-time setting, but several transfer to continuous-time. As well as the mixture process, some of our results also cover the pure exclusion case. We state several significant open problems that remain.

Key words and phrases: Exclusion process; voter model; passage-time moments; hybrid zone; almost-sure bounds; Lyapunov functions.

AMS 2000 Mathematics Subject Classification: 60K35 (Primary); 60J05, 60J25, 82C22 (Secondary).

1 Introduction

The *exclusion-voter* model studied in this paper is a one-dimensional lattice-based interacting-particle process with nearest-neighbour interactions, introduced by Belitsky *et al.* in [5], that is a mixture of the symmetric voter model and the simple exclusion process. For background on the exclusion process and voter model (separately), and interacting particle systems in general, see e.g. [14, 15].

*Partially supported by the Heilbronn Institute for Mathematical Research

The voter model has been used to model for the spread of an opinion, such as voting intention, through a static population via nearest-neighbour interactions; see e.g. [11]. The mixture model studied here is a natural extension of this model, whereby individual particles (voters) do not have to remain static, but may move by switching places. Alternative motivation, such as from the point of view of competition of species (see e.g. [6]) also can be adapted to the mixture model. As our results show, allowing place-swaps can have a dramatic effect on the dynamics of the process.

The exclusion-voter model is a Markov process with state-space $\{0,1\}^{\mathbb{Z}}$; each site of the one-dimensional lattice $\mathbb{Z}$ can be labelled either 0 or 1, where the labels 0, 1 represent two types of particle. The ground-state of our model will be the ‘Heaviside’ configuration $\dots 111000 \dots$. We will consider initial configurations that are finite perturbations of this ground-state, and so will contain a finite number of unlike pairs, where by ‘pair’ we always mean two adjacent particles.

In this paper, we concentrate on a *discrete time* process that can be described informally as follows. At each time step, the *simple exclusion process* selects uniformly at random from amongst all unlike pairs. If the chosen pair is 01, it flips to 10 with probability p (else there is no change); if the pair is 10, it flips to 01 with probability $1 - p$. On the other hand, at each time step the *symmetric voter model* selects uniformly at random from all unlike pairs and then flips the chosen pair to either 00 or 11, with equal chance of each. The model that is considered in this paper, introduced in [5], is a mixture of these two processes, whereby at each time step we determine independently at random whether to perform a voter-type move (with probability β) or an exclusion-type move (probability $1 - \beta$).

The analogous continuous-time exclusion-voter model can be defined via its infinitesimal generator and constructed via a Harris-type graphical construction. Then the discrete-time process described above is naturally embedded in the continuous-time process. In our analysis we work in discrete-time, and the discrete-time process has its own interest, but, as we shall indicate, some of our results transfer almost immediately into continuous time.

Individually, the exclusion process and voter model exhibit very different behaviour. For instance, in the exclusion process there is local conservation of 1s: the number of 1s in a bounded interval can change only through the boundary. There is no such conservation in the voter model. In the mixture process that we study in the present paper, voter moves and exclusion moves interact in a highly non-trivial way. This introduces technical difficulties: for instance, voter moves can cause drastic changes quickly, and there is no obvious monotonicity property (as mentioned in [5]). We describe the model more formally and state our results in the next section. First we outline the existing literature and the contribution of the present paper.

In [5], the exclusion-voter model was introduced, and results were proved for the exclusion process and voter model separately, as well as some initial results for the mixture model. The main problems left open in [5] were the non-existence of passage-time moments and the issue of transience/recurrence for the mixture model. As we describe shortly, the present paper makes contributions to each of these problems. Some of the results in [5], in the symmetric exclusion ($p = 1/2$) case, are generalized to non-nearest-neighbour interactions in [20]. Certain ‘ergodic’ properties of a generalization of the continuous-time exclusion-voter model, again in the symmetric exclusion case, are studied in [12]; the context of that paper is different from that of the present paper or [5]. The goal of the present paper is to study

the mixture model in more depth than [5]. In particular we prove new results on: (i) the passage-time problem for the exclusion-voter model, the main contribution being the (more difficult) non-existence of passage-time moments for the mixture model; and (ii) the size of the disordered region where 1s and 0s intermingle. This disordered region we call the *hybrid zone* (cf. [7]). Our results still leave several interesting open problems, and we make some conjectures with regard to these in the next section.

As described above, one contribution of the present paper is to address the passage-time problem for the exclusion-voter model. The passage-time of interest to us here is the *relaxation time* — the return time of the configuration to the ground-state. Let us briefly outline the passage-time problem in some generality. On a suitable probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with expectation $\mathbb{E}$, let $(X_t)_{t \in \mathbb{Z}^+}$ be a discrete-time stochastic process, with state-space $\mathcal{S}$. For $A \subseteq \mathcal{S}$ set

$$\tau := \min\{t \in \mathbb{N} : X_t \in A\}.$$

Here and throughout the paper we adopt the conventions that $\mathbb{Z}^+ := \{0, 1, 2, \dots\}$, $\mathbb{N} := \{1, 2, 3, \dots\}$, and $\min \emptyset := +\infty$. The stopping time τ is the *first passage time* of A .

A fundamental question is whether τ is almost-surely finite. If so, one is interested in whether $\mathbb{E}[\tau]$ is finite, or, more generally, whether for $s > 0$ the (generalized) moment $\mathbb{E}[\tau^s]$ is finite. In general, often one can prove the existence of moments directly via semimartingale (Lyapunov-type function) criteria such as those in [2, 4, 13] in the vein of Foster [10].

The non-existence of moments (for which no results were previously obtained for the exclusion-voter model with $\beta \in (0, 1)$) is usually a harder problem. In general, semimartingale-type arguments are available in this case too (see e.g. [3, 4, 13]), but under more restrictive conditions than the corresponding existence results: non-existence results typically need fine control over jumps of the process. Lamperti [13] was first to establish a general methodology for proving non-existence of passage-time moments, based upon finding a suitable submartingale and obtaining a good-probability lower bound for passage times; his method was later extended in [3, 4]. The same two elements form the basis of our approach, but we must proceed differently since the exclusion-voter model does not possess the regularity required by existing general results such as those of [3, 4, 13].

Let us describe more specifically the contribution of the present paper to the passage-time problem for the exclusion-voter model. On the one hand, we extend the region of the parameter space of the model for which almost-sure finiteness of the relaxation time is known, and we also obtain results on the existence of higher moments of the relaxation time (including in the case of pure exclusion). On the other hand, we show the *non-existence* of certain passage-time moments; this problem was not addressed in [5]. Each of these opposed directions requires us to develop new techniques.

In the present paper we prove, for example, that under certain conditions $1 + \varepsilon$ moments ($\varepsilon > 0$) of the passage-time do not exist; this approaches an outstanding conjecture in [5]. Although the conjecture in [5] remains open, by developing methods to prove non-existence of passage-time moments in this context, this paper makes a significant step towards settling that conjecture. The argument for non-existence of moments in the present paper will rest crucially on a good-probability lower bound (Lemma 9 below) for the time taken for the process to reach the target state; we comment more fully on this at the appropriate point in our argument.

The second main contribution of the paper is to address the evolution of the size of the hybrid zone where 0s and 1s intermingle. Here our basic tools are again semimartingales: we apply some general results on obtaining almost-sure bounds for stochastic processes via semimartingale-type criteria from [17]. For instance, for the pure exclusion process in the case $p = 1/2$ we prove that with probability 1 the maximum size of the hybrid zone up to time t remains bounded between $t^{1/3}$ and $t^{1/2}$, ignoring logarithmic factors.

There are various connections between the exclusion-voter model and other stochastic models. The model that we consider fits into the general family of two-species reaction-diffusion processes, many other examples of which have been studied; see e.g. [18]. The model can be reinterpreted as a growth/depletion model for a randomly evolving domain or interface (see Section 4.2 below). Also, the pure voter model has connections to zero-drift random walks in the quarter-plane (see Section 6 of [5]); the addition of the exclusion process can be viewed as a perturbation of this random walk model. Thus the exclusion-voter model has features in common with perturbed random walks (see e.g. [16]).

In some cases these connections to other processes are particularly fruitful. The exclusion process in either of the extreme cases $p \in \{0, 1\}$ is known as *totally asymmetric*. At least in the continuous-time version of the process, there is a now well-known relation between the totally asymmetric simple exclusion process (TASEP) in 1 dimension and last-passage percolation in 2 dimensions and related combinatorial models; see e.g. [9]. As a result, TASEP is now rather well understood. However, this useful connection is not available for the general exclusion-voter model, which has much more complicated dynamics.

In the next section we give some more formal definitions and state our main results for the exclusion-voter model. There remain many open problems for this model, and we draw attention to some of these as we go. We believe that these problems are challenging but that further progress is possible towards their solution by building upon the techniques in the present paper.

2 Definitions and statement of results

We now formally describe the model that we study, as considered in [5]. We introduce some notation to describe the configuration of the process. Let $\mathcal{D}' \subset \{0, 1\}^{\mathbb{Z}}$ denote the set of configurations with a finite number of 0s to the left of the origin and 1s to the right. Let ' $\sim$ ' denote the equivalence relation on $\mathcal{D}'$ such that for $S, S' \in \mathcal{D}'$, $S \sim S'$ if and only if S and S' are translates of each other. Then set $\mathcal{D} := \mathcal{D}' / \sim$. In other words, the configuration space $\mathcal{D}$ is the set of configurations of the form infinite string of 1s—finite number of 0s and 1s—infinite string of 0s, modulo translations. For example, one configuration $S \in \mathcal{D}$ is

$$S = \dots 1110000111001010100001111000 \dots \quad (1)$$

Configurations such as those in $\mathcal{D}$ are sometimes called *shock profiles* (see e.g. [5] and references therein).

Fix $\beta \in [0, 1]$ (the mixing parameter) and $p \in [0, 1]$ (the exclusion parameter). The discrete-time exclusion-voter process $\xi = (\xi_t)_{t \in \mathbb{Z}^+}$ with parameters (β, p) is a time-homogeneous Markov chain on the countable state-space $\mathcal{D}$. The one-step transition probabilities are determined by the following mechanism. At each time step we decide independently at random whether to perform a *voter* move or an *exclusion* move. We choose a voter

move with probability β and an exclusion move with probability $1 - \beta$. Having decided this, choose an unlike adjacent pair (i.e. 01 or 10) uniformly at random from all possibilities on which to perform the move.

The voter move is such that the chosen pair (01 or 10) flips to 00 or 11 each with probability $1/2$. The exclusion move is such that a chosen pair 01 flips to 10 with probability p (otherwise no move) and a chosen pair 10 flips to 01 with probability $q := 1 - p$ (otherwise no move).

In addition to the discrete-time model that is the focus of the present paper, there is a corresponding continuous-time model, also introduced in [5]. A priori, the relationship between the two time-scales is complicated, but from our results on the discrete-time process we can obtain some results in the continuous-time setting too. For a description of the continuous-time model, its relationship to the discrete-time model that is our main object of study, and our results in that case, see Section 3 below.

The underlying probability space for ξ we denote by $(\Omega, \mathcal{F}, \mathbb{P}_{\beta,p})$, and the corresponding expectation $\mathbb{E}_{\beta,p}$. We denote the ground-state Heaviside configuration $\mathcal{D}_0 \in \mathcal{D}$, which consists of a single pair 10 abutted by infinite strings of 1s and 0s to the left and right, respectively. That is,

$$\mathcal{D}_0 = \dots 11110000 \dots,$$

up to translation. The next result gives some elementary properties of the state-space $\mathcal{D}$ under $\mathbb{P}_{\beta,p}$. In particular, Proposition 1 says that for $(\beta, p) \in (0, 1)^2$ (i.e., in the interior of the parameter space) ξ is irreducible and aperiodic under $\mathbb{P}_{\beta,p}$.

Proposition 1 *$\mathcal{D}_0$ is an absorbing state under $\mathbb{P}_{\beta,1}$ for any $\beta \in [0, 1]$. Suppose $\beta \neq 1$ and $(\beta, p) \notin \{(0, 0), (0, 1)\}$. Then all states in $\mathcal{D} \setminus \{\mathcal{D}_0\}$ communicate under $\mathbb{P}_{\beta,p}$. Suppose $\beta \neq 1$, $p < 1$, and $(\beta, p) \neq (0, 0)$. Then all states in $\mathcal{D}$ communicate under $\mathbb{P}_{\beta,p}$, and ξ is irreducible and aperiodic.*

For $S_0 \in \mathcal{D}$ define the hitting time of $\mathcal{D}_0$ for the process ξ as

$$\tau := \min\{t \in \mathbb{N} : \xi_t = \mathcal{D}_0\}.$$

We call τ the *relaxation time* of the system. We introduce some convenient terminology. If $\mathbb{P}_{\beta,p}(\tau = +\infty \mid \xi_0 = S_0) > 0$ for $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$, we say that ξ is *transient* started from S_0 ; if $\mathbb{P}_{\beta,p}(\tau < \infty \mid \xi_0 = S_0) = 1$ for $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$, we say that ξ is *recurrent* started from S_0 . In the latter case, if in addition $\mathbb{E}_{\beta,p}[\tau \mid \xi_0 = S_0] < \infty$ for $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$, we say that ξ is *positive-recurrent* started from S_0 . When ξ is irreducible (see Proposition 1), this terminology coincides with the standard usage for countable state-space Markov chains. Further, when ξ is irreducible and aperiodic (see Proposition 1), we may use the term *ergodic* in the positive-recurrent case.

Results of Liggett (see e.g. Chapter VIII of [14]) imply that the pure exclusion process ($\beta = 0$) is positive-recurrent for all $S_0 \in \mathcal{D}$ if and only if $p > 1/2$. We recall the following result, which is contained in Theorems 5.1, 5.2, 6.1, 7.1, and 7.2 of [5], together with an inspection of (7.2) in [5] for part (iii)(a).

Theorem 1 *(i) Suppose $\beta = 0$ (pure exclusion). Then for any $S_0 \in \mathcal{D}$, ξ is positive-recurrent for $p > 1/2$ and transient for $p \leq 1/2$.*

(ii) Suppose $\beta = 1$ (pure voter). Then ξ is positive-recurrent for any $S_0 \in \mathcal{D}$, and moreover for any $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ and any $\varepsilon > 0$,

$$\mathbb{E}_{1,p}[\tau^{(3/2)-\varepsilon} \mid \xi_0 = S_0] < \infty; \quad \mathbb{E}_{1,p}[\tau^{(3/2)+\varepsilon} \mid \xi_0 = S_0] = \infty.$$

(iii) Suppose $\beta \in (0, 1)$ (mixture process). Then

(a) If β and $p \in [0, 1]$ are such that $(1-p)(1-\beta) < 1/3$, then ξ is positive-recurrent for any $S_0 \in \mathcal{D}$. In particular, for any $\beta > 2/3$ and any $p \in [0, 1]$, ξ is positive-recurrent for any $S_0 \in \mathcal{D}$.

(b) For $p \geq 1/2$ and any $\beta > 0$, ξ is positive-recurrent for any $S_0 \in \mathcal{D}$.

In [5], the following was Conjecture 7.1.

Conjecture 1 For any $p < 1/2$, there exists $\beta_0 = \beta_0(p) > 0$ such that for all $\beta < \beta_0$, ξ is not positive-recurrent, i.e. $\mathbb{E}_{\beta,p}[\tau \mid \xi_0 = S_0] = \infty$ for any $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$.

Our first result says that for β small enough (so that the exclusion part is prevalent), $1 + \varepsilon$ moments do not exist; thus Conjecture 1 remains tantalizingly open.

Theorem 2 For each $p < 1/2$, there exists a constant $\beta_1 = \beta_1(p) = (1 - 2p)/(2 - 2p) \in (0, 1/2]$ such that for all $\beta \leq \beta_1$, for any $\varepsilon > 0$, for any $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$,

$$\mathbb{E}_{\beta,p}[\tau^{1+\varepsilon} \mid \xi_0 = S_0] = \infty.$$

Our second result (which is immediate for $\beta = 0$, by Theorem 1(i), and for $\beta = 1$, by Theorem 1(ii)) says that in the mixture process, the presence of a transient exclusion ensures that $2 + \varepsilon$ moments do not exist. Thus for $p \leq 1/2$, even in the case where Theorem 1(iii) applies the recurrence is polynomial in nature, i.e. ‘heavy-tailed’.

Theorem 3 Suppose that $p \leq 1/2$ and $\beta \in [0, 1]$. Then for any $\varepsilon > 0$ and $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$,

$$\mathbb{E}_{\beta,p}[\tau^{2+\varepsilon} \mid \xi_0 = S_0] = \infty.$$

In view of Theorem 1(ii), we suspect that mixing transient ($p \leq 1/2$) exclusion with the voter model should not lead to finiteness of more moments for τ , and so we make the following conjecture.

Conjecture 2 Suppose that $p \leq 1/2$ and $\beta \in [0, 1]$. Then for any $\varepsilon > 0$ and $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$,

$$\mathbb{E}_{\beta,p}[\tau^{(3/2)+\varepsilon} \mid \xi_0 = S_0] = \infty.$$

Even this conjecture seems to be challenging. The obstacle is that the exclusion and voter moves can interact in complex ways. Technically, the issue that prevents us from reducing the 2 to 3/2 in Theorem 3 is that the exclusion process can (and typically will) increase the number of blocks. Intuitively, the behaviour of the system is very unclear.

As a particular example of our theorems above, suppose $\beta > 0$, $p = 1/2$, and $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$. Then we have from Theorem 1(iii)(b) that $\mathbb{E}_{\beta,1/2}[\tau \mid \xi_0 = S_0] < \infty$ (indeed,

$\mathbb{E}_{\beta,1/2}[\tau^{1+\varepsilon} \mid \xi_0 = S_0] < \infty$ for some $\varepsilon = \varepsilon(\beta) > 0$ by Remark 7.2 in [5]) while on the other hand Theorem 3 says that $\mathbb{E}_{\beta,1/2}[\tau^{2+\varepsilon} \mid \xi_0 = S_0] = \infty$ for any $\varepsilon > 0$.

An open problem mentioned in [5] is whether the mixture process with $\beta > 0$ and $p < 1/2$ is in fact transient (it is recurrent for $p \geq 1/2$, by Theorem 1(iii)(b)). Simulations that we have performed have been inconclusive. We conjecture the following.

Conjecture 3 *Suppose that $p < 1/2$ and $\beta > 0$. Then for any $S_0 \in \mathcal{D}$, ξ is recurrent, i.e., $\mathbb{P}_{\beta,p}(\tau < \infty \mid \xi_0 = S_0) = 1$.*

Let us indicate, very roughly, some motivation behind Conjecture 3. As demonstrated in [5] (Section 6) the two-block voter model can be mapped into a zero-drift random walk in the positive quadrant; the relaxation time of the system is then the exit-time of the walk from the quadrant. Mixing in the exclusion process may be viewed as adding a small perturbation to this walk. Analogy with exit-from-cones problems for perturbed random walks in [16] suggests that this exit time remains almost-surely finite.

Note that if Conjectures 1 and 3 both hold, there is *null-recurrence* for $p < 1/2$ and $\beta \in (0, \beta_0)$. Our next result represents some progress in the direction of Conjecture 3, and gives recurrence in a previously unexplored region of the parameter space.

Theorem 4 *Suppose that $p < 1/2$ and $\beta \geq 4/7$. Then for any $S_0 \in \mathcal{D}$, ξ is recurrent, i.e., $\mathbb{P}_{\beta,p}(\tau < \infty \mid \xi_0 = S_0) = 1$.*

Now we turn to the problem of existence of moments for τ . Our first result deals with the pure exclusion process in the positive-recurrent ($p > 1/2$) case. If we further restrict to $p > 2/3$, it is possible to construct a positive strict supermartingale with uniformly bounded increments (see (5.7) in [5]), and so it is not hard to show that all polynomial moments of τ exist in that case. Theorem 5 below extends this conclusion to all $p > 1/2$.

Theorem 5 *Suppose that $\beta = 0$ and $p > 1/2$. Then for any $S_0 \in \mathcal{D}$ and any $s \in [0, \infty)$*

$$\mathbb{E}_{0,p}[\tau^s \mid \xi_0 = S_0] < \infty.$$

We suspect that, under the conditions of Theorem 5, the existence of some super-polynomial ‘moments’ for τ can be obtained via our techniques and general results from [3]. The next result covers the mixture process in the case where the exclusion component is positive-recurrent. In the $\beta \in [0, 1]$, $p > 1/2$ case we know from Theorem 1 that $\mathbb{E}[\tau] < \infty$; the next theorem says that some higher moments are finite also.

Theorem 6 *Suppose that $\beta \in [0, 1]$ and $p > 1/2$. Then for any $S_0 \in \mathcal{D}$*

$$\mathbb{E}_{\beta,p}[\tau^{6/5} \mid \xi_0 = S_0] < \infty.$$

In view of Theorem 1 and Theorem 5, in the setting of Theorem 6 we are mixing together the voter model, for which $(3/2) - \varepsilon$ moments exist, and the recurrent exclusion process, for which all moments exist. Thus one might hope to improve the exponent in Theorem 6 to at least $(3/2) - \varepsilon$, although it is again not at all clear whether the two processes interact in this way.




Figure 1: Representations of the (β, p) parameter space. The key given explains the labelling together with the appropriate result from the text (for brevity we have dropped the subscripts on $\mathbb{P}, \mathbb{E}$ in the table).

Figure 1 gives two diagrams of the (β, p) parameter space, summarizing the results of the previous theorems for the relaxation time τ .

We now state our results on the size of the hybrid zone. First we need to introduce some more notation, following [5]. A 1-block (0-block) is a maximal string of consecutive 1s (0s). Configurations in $\mathcal{D}$ consist of a finite number of such blocks. For $S \in \mathcal{D}$, let $N = N(S) \geq 0$ denote the number of 1-blocks not including the infinite 1-block to the left (this is the same as number of 0-blocks not including the infinite 0-block to the right). Enumerating left to right, let $n_i = n_i(S)$ denote the size of the i -th 0-block, and $m_i = m_i(S)$ the size of the i -th 1-block. Then we may represent configuration $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ by the vector $(n_1, m_1, \dots, n_N, m_N)$. For example, the configuration S of (1), which has $N(S) = 5$, has the representation $(4, 3, 2, 1, 1, 1, 1, 1, 4, 4)$.

Set $|\mathcal{D}_0| := 0$ and for $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ let $|S| := \sum_{i=1}^N (n_i + m_i)$ the size of the hybrid zone, i.e., the length of the string of 0s and 1s between the infinite string of 1s to the left and the infinite string of 0s to the right.

The next result gives upper bounds for the size of the hybrid zone $|\xi_t|$ and the number of blocks $N(\xi_t)$; in particular, part (ii) covers the case $\beta = 0, p = 1/2$ of the symmetric pure (transient) exclusion process. (Also included is the pure voter model $\beta = 1$, but in that case $\mathcal{D}_0$ is absorbing and ξ positive-recurrent, so that $\mathbb{P}_{1,p}(\lim_{t \rightarrow \infty} |\xi_t| = 0) = 1$.)

Theorem 7 (i) Suppose $\beta \in [0, 1], p \in [0, 1]$. For any $\varepsilon > 0$, $\mathbb{P}_{\beta,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$,

$$\max_{0 \leq s \leq t} N(\xi_s) \leq \begin{cases} t^{1/2} (\log t)^{(1/2)+\varepsilon} & \text{if } p < 1/2 \\ t^{1/3} (\log t)^{(1/3)+\varepsilon} & \text{if } p \geq 1/2 \end{cases} \quad (2)$$

(ii) Suppose $\beta \in [0, 1]$, $p \geq 1/2$. For any $\varepsilon > 0$, $\mathbb{P}_{\beta,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$,

$$\max_{0 \leq s \leq t} |\xi_s| \leq t^{1/2}(\log t)^{(1/2)+\varepsilon}. \quad (3)$$

The remainder of our results deal with the pure exclusion process ($\beta = 0$). In the continuous-time setting, related results on the growth of the hybrid zone of the pure exclusion process were first obtained by Rost [19] (in the totally asymmetric case); see Section VIII.5 of [14], and [1] for more general results. In particular, Theorems 5.2 and 5.2 of [14] say, very loosely, that under $\mathbb{P}_{0,p}$

$$|\eta_t| \approx t \quad (p < 1/2); \quad |\eta_t| \approx t^{1/2} \quad (p = 1/2),$$

where η is the continuous-time version of ξ as described in Section 3. In particular, the symmetric case is significantly different from the asymmetric case. However, there seems to be no immediate way to translate these results between the continuous- and discrete-time settings (see Section 3 below).

Part (i) of the next result strengthens the bound in (2) slightly in the pure exclusion case with $p < 1/2$. Part (ii) complements the $\beta = 0$ case of (3) for the case $p < 1/2$ (transient but not symmetric exclusion); in particular it quantifies the rate of transience.

Theorem 8 *Suppose $\beta = 0$ and $p \in [0, 1]$.*

(i) *There exists $C \in (0, \infty)$ such that for any $p \in [0, 1]$, $\mathbb{P}_{0,p}$ -a.s., for all $t \in \mathbb{Z}^+$,*

$$\max_{0 \leq s \leq t} N(\xi_s) \leq Ct^{1/2}.$$

(ii) *Suppose $p \in [0, 1/2)$. Then for any $\varepsilon > 0$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$*

$$\max_{0 \leq s \leq t} |\xi_s| \leq t^{2/3}(\log t)^{(1/3)+\varepsilon}.$$

On the other hand, there exists $c(p) \in (0, \infty)$ such that for any $c \in (0, c(p))$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$

$$|\xi_t| \geq ct^{1/2}.$$

Our next result complements (3) in the case $\beta = 0$, $p = 1/2$. Information on $N(\xi_t)$ in this case is given by the $p = 1/2$ case of (2).

Theorem 9 *Suppose $\beta = 0$, $p = 1/2$. For any $\varepsilon > 0$, $\mathbb{P}_{0,1/2}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$,*

$$t^{1/3}(\log t)^{-(1/3)-\varepsilon} \leq \max_{0 \leq s \leq t} |\xi_s| \leq t^{1/2}(\log t)^{(1/2)+\varepsilon}.$$

It is an open problem to obtain sharper versions of the above results on $|\xi_t|$. In the pure exclusion ($\beta = 0$) case, we conjecture the following.

	$0 \leq \beta \leq 1$	$\beta = 0$
$p < 1/2$	?	$t^{2/3}(\log t)^{(1/3)+\varepsilon}$
$p \geq 1/2$	$t^{1/2}(\log t)^{(1/2)+\varepsilon}$	

Table 1: Upper bound for $\max_{0 \leq s \leq t} |\xi_s|$.

	$0 \leq \beta \leq 1$	$\beta = 0$
$p < 1/2$	$t^{1/2}(\log t)^{(1/2)+\varepsilon}$	$Ct^{1/2}$
$p \geq 1/2$	$t^{1/3}(\log t)^{(1/3)+\varepsilon}$	

Table 2: Upper bound for $\max_{0 \leq s \leq t} N(\xi_s)$.

Conjecture 4 Suppose $\beta = 0$. If $p < 1/2$, then for any $\varepsilon > 0$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$

$$|\xi_t| \leq t^{(1/2)+\varepsilon}.$$

If $p = 1/2$, then for any $\varepsilon > 0$, $\mathbb{P}_{0,1/2}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$

$$t^{(1/3)-\varepsilon} \leq |\xi_t| \leq t^{(1/3)+\varepsilon}.$$

We summarize some of our results on $|\xi_t|$, $N(\xi_t)$ in Tables 1 and 2.

The structure of the remainder of the paper is as follows. In Section 3 we describe the *continuous time* version of the exclusion-voter model, and how it relates to the discrete-time version studied here, and which results can be transferred without too much extra work. Section 4 collects some preliminary results: in Section 4.1 we collect general semimartingale results that we apply in the paper, in Section 4.2 we introduce notation and a convenient representation for configurations of the model and give a proof of Proposition 1, and in Section 4.3 we give some lemmas on the Lyapunov-type functions that we will use throughout the paper. In Section 5 we present the proofs of Theorems 2 and 3 on passage-time moments, via a series of lemmas. In Section 6 we prove Theorem 4. In Section 7 we prove Theorems 5 and 6. In Section 8 we give the proofs of Theorems 7, 8, and 9 on the size of the hybrid zone and number of blocks.

3 Continuous time

The exclusion process and voter model are often studied in continuous time [14, 15]. Their mixture, the exclusion-voter model, may also be defined in continuous time, as in [5]. We recall the definition, following [5].

Let $\nu = (\nu(x))_{x \in \mathbb{Z}} \in \{0, 1\}^{\mathbb{Z}}$, so that $\nu(x)$ is the label (0 or 1) at x . For $x, y, z \in \mathbb{Z}$ denote

$$\nu_{x,y}(z) := \begin{cases} \nu(y) & \text{if } z = x, \\ \nu(x) & \text{if } z = y, \\ \nu(z) & \text{if } z \neq x, y; \end{cases} \quad \nu_x(z) := \begin{cases} 1 - \nu(z) & \text{if } z = x, \\ \nu(z) & \text{if } z \neq x. \end{cases}$$

In words, $\nu_{x,y}$ is ν with labels at x, y interchanged, and ν_x is ν with the label at x flipped (i.e., replaced with its opposite).

We introduce Markovian generators Ω_p^e ($p \in [0, 1]$) and Ω^v defined by their action on functions f on $\{0, 1\}^{\mathbb{Z}}$ by:

$$\Omega_p^e f(\nu) = \sum_{x,y} p(x,y) \nu(x) (1 - \nu(y)) [f(\nu_{x,y}) - f(\nu)],$$

where

$$p(x,y) := \begin{cases} p & \text{if } y = x - 1, \\ 1 - p & \text{if } y = x + 1, \\ 0 & \text{otherwise;} \end{cases}$$

and by

$$\Omega^v f(\nu) = \sum_x c(x, \nu) [f(\nu_x) - f(\nu)],$$

where

$$c(x, \nu) := \begin{cases} \frac{1}{2} (\nu(x-1) + \nu(x+1)) & \text{if } \nu(x) = 0, \\ \frac{1}{2} (2 - \nu(x-1) - \nu(x+1)) & \text{if } \nu(x) = 1. \end{cases}$$

The continuous-time simple exclusion process with parameter p has generator Ω_p^e . Informally, a 1 at site x switches places with a 0 at $x + 1$ at rate $1 - p$ and with a 0 at site $x - 1$ with rate p . The continuous-time voter model has generator Ω^v . Informally, a particle (0 or 1) at site x flips to a particle of its opposite kind at rate proportional to the number of neighbours of x occupied by particles of its opposite kind.

The continuous-time exclusion-voter model with mixing parameter $\beta \in [0, 1]$ and exclusion parameter $p \in [0, 1]$ is a Markov process $(\eta'_t)_{t \geq 0}$ on $\mathcal{D}' \subset \{0, 1\}^{\mathbb{Z}}$ with generator

$$(1 - \beta)\Omega_p^e + \beta\Omega^v.$$

This induces a Markov process $\eta = (\eta_t)_{t \geq 0}$ on the space of equivalence classes $\mathcal{D}$ by taking η_t to be the $\sim$ -equivalence class of η'_t .

The process η can be constructed from an array of homogeneous one-dimensional Poisson processes via a Harris-type graphical construction, which we briefly (and rather informally) review: see p. 9 of [5]. Independently for each $x \in \mathbb{Z}$, we construct four independent Poisson processes on $[0, \infty)$, $(\mathcal{N}_t^{x,x+1})_{t \geq 0}$, $(\mathcal{N}_t^{x,x-1})_{t \geq 0}$, $(\mathcal{M}_t^{x,x+1})_{t \geq 0}$, $(\mathcal{M}_t^{x,x-1})_{t \geq 0}$, with rates $(1 - \beta)p$, $(1 - \beta)(1 - p)$, $\beta/2$, $\beta/2$ respectively. Given the initial configuration η_0 , the dynamics of η is then constructed as follows. If there is a Poisson arrival at time t in $\mathcal{N}^{x,x+1}$ (resp., $\mathcal{N}^{x,x-1}$), and if at time t^- $x = 1$ while $x + 1 = 0$ (resp., $x - 1 = 0$), then the 1 at x switches with the 0 at $x + 1$ (resp., $x - 1$) at time t . If there is a Poisson arrival at time t in $\mathcal{M}^{x,x+1}$ (resp., $\mathcal{M}^{x,x-1}$), then the site $x + 1$ (resp., $x - 1$) acquires the same state at time t as the state of x at time t^- .

Let $\mu_0 := 0$ and for $n \in \mathbb{N}$ set

$$\mu_n := \inf \left\{ t > \mu_{n-1} : \sum_{x,y:|x-y|=1} |\eta_{t^-}(x) - \eta_{t^-}(y)| \times (\mathcal{N}^{x,y}(\mu_{n-1}, t] + \mathcal{M}^{x,y}(\mu_{n-1}, t]) > 0 \right\},$$

where $\mathcal{N}(s, t]$ denotes the number of Poisson arrivals in $(s, t]$ for the process $\mathcal{N}$. The times μ_n are then the instants of *attempted jumps* of η . It is now not hard to see that with the definitions in Section 2 and this section, ξ may be embedded in η via

$$(\xi_t)_{t \in \mathbb{Z}^+} \stackrel{d}{=} (\eta_{\mu_t})_{t \in \mathbb{Z}^+},$$

given $\xi_0 = \eta_0 = S_0 \in \mathcal{D}$, where ‘ $\stackrel{d}{=}$ ’ stands for equality in distribution.

In the continuous-time setting, the relaxation time is

$$\tau_c := \inf\{t \geq 0 : \eta_t = \mathcal{D}_0\}.$$

The natural question is: given the results in Section 2 on τ , what is it possible to say about τ_c ? We now outline which of our discrete-time results for ξ can be readily transferred to continuous-time results for η . Compare Section 8 of [5].

First of all, as pointed out in [5], recurrence and transience transfer directly:

$$\mathbb{P}_{\beta,p}(\tau < \infty \mid \xi_0 = S_0) = 1 \iff \mathbb{P}_{\beta,p}(\tau_c < \infty \mid \eta_0 = S_0).$$

To conclude about moments (i.e., tails) of the relaxation times, rather than merely their finiteness, it is necessary to know about the comparative rates of the two processes. Observe that the transition rate of the continuous-time process is, roughly speaking, proportional to the number of blocks: the continuous-time process tends to evolve at least as fast as the discrete-time process.

The pure voter model ($\beta = 1$) is well-behaved in the sense that it cannot increase the number of blocks. Thus, roughly speaking, the discrete and continuous timescales are directly comparable, and results are more easily transferred. This intuition is formalized in Section 8 of [5], where it is shown that for any $s > 0$

$$\mathbb{E}_{1,p}[\tau^s \mid \xi_0 = S_0] < \infty \iff \mathbb{E}_{1,p}[\tau_c^s \mid \eta_0 = S_0] < \infty.$$

In the more general mixture setting, without more information about the evolution of the number of blocks, only one-sided results are possible a priori. In particular, it is shown in Section 8 of [5] that for any $s > 0$

$$\mathbb{E}_{\beta,p}[\tau^s \mid \xi_0 = S_0] < \infty \implies \mathbb{E}_{\beta,p}[\tau_c^s \mid \eta_0 = S_0] < \infty.$$

So Theorem 1 due to [5] transfers directly to continuous time, and holds with τ_c instead of τ ; this is Theorem 1.1 in [5]. Moreover our Theorems 4, 5 and 6 also carry across, and hold with τ_c . Thus we obtain the following corollaries.

Corollary 1 *Suppose that $p < 1/2$ and $\beta \geq 4/7$. Then for any $S_0 \in \mathcal{D}$, η is recurrent, i.e., $\mathbb{P}_{\beta,p}(\tau_c < \infty \mid \eta_0 = S_0) = 1$.*

Corollary 2 *Suppose that $\beta = 0$ and $p > 1/2$. Then for any $S_0 \in \mathcal{D}$ and any $s \in [0, \infty)$*

$$\mathbb{E}_{0,p}[\tau_c^s \mid \xi_0 = S_0] < \infty.$$

Corollary 3 *Suppose that $\beta \in [0, 1]$ and $p > 1/2$. Then for any $S_0 \in \mathcal{D}$*

$$\mathbb{E}_{\beta,p}[\tau_c^{6/5} \mid \xi_0 = S_0] < \infty.$$

In particular, Corollary 2 says that for the standard (continuous-time) recurrent exclusion process, all moments of τ_c exist. This fact may well be known, but we could not find a reference in the literature. However, the general non-existence of moments and size of the hybrid-zone results do not carry across by these existing arguments. With some extra technical work, it is likely that some of the other results given in Section 2 can be translated in some form to continuous time. We do not address this issue here.

4 Preliminaries

4.1 Technical tools

In this section we state some general martingale-type results that we will need. In particular, we will recall some criteria for obtaining upper and lower almost sure bounds for discrete-time stochastic processes on the half-line given in [17]. We will apply these results to the hybrid zone of the exclusion-voter model in Section 8 below.

Let $(\mathcal{F}_t)_{t \in \mathbb{Z}^+}$ be a filtration on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $X = (X_t)_{t \in \mathbb{Z}^+}$ be a discrete time $(\mathcal{F}_t)$ -adapted stochastic process taking values in $[0, \infty)$. Suppose that $\mathbb{P}(X_0 = x_0) = 1$ for some $x_0 \in [0, \infty)$. For the applications in the present paper, we will for instance take $X_t = |\xi_t|$.

We will need the following maximal inequality:

Lemma 1 *Let $(X_t)_{t \in \mathbb{Z}^+}$ be an $(\mathcal{F}_t)$ -adapted process on $[0, \infty)$ such that $\mathbb{P}(X_0 = x_0 \mid \mathcal{F}_0) = 1$ and for some $B \in (0, \infty)$ and all $t \in \mathbb{Z}^+$*

$$\mathbb{E}[X_{t+1} - X_t \mid \mathcal{F}_t] \leq B \text{ a.s..} \quad (4)$$

Then for any $r > 0$ and any $t \in \mathbb{N}$

$$\mathbb{P}\left(\max_{0 \leq s \leq t} X_s \geq r\right) \leq (Bt + x_0)r^{-1}. \quad (5)$$

Proof. Similarly to Doob's decomposition (see e.g. [21], p. 120), set $Y_0 := X_0$, and for $s \in \mathbb{N}$ let $Y_s := X_s + A_{s-1}^- + A_{s-2}^- + \cdots + A_1^-$, where $A_s = \mathbb{E}[X_{s+1} - X_s \mid \mathcal{F}_s]$, $A_s^- = \max\{-A_s, 0\} \geq 0$, $A_s^+ = \max\{A_s, 0\} \geq 0$. Then

$$\mathbb{E}[Y_{s+1} - Y_s \mid \mathcal{F}_s] = \mathbb{E}[X_{s+1} - X_s + A_s^- \mid \mathcal{F}_s] = A_s + A_s^- = A_s^+ \in [0, B]$$

so that Y_s is a nonnegative $(\mathcal{F}_s)$ -submartingale with $Y_s \geq X_s$ for all s , and $\mathbb{E}[Y_t] \leq Y_0 + Bt = x_0 + Bt$. Consequently, by Doob's submartingale inequality (see e.g. [21], p. 137)

$$\mathbb{P}\left(\max_{0 \leq s \leq t} X_s \geq r\right) \leq \mathbb{P}\left(\max_{0 \leq s \leq t} Y_s \geq r\right) \leq r^{-1} \mathbb{E}[Y_t] \leq (Bt + x_0)r^{-1},$$

as required. $\square$

The next result is contained in Theorem 3.2 of [17].

Lemma 2 *Let $(X_t)_{t \in \mathbb{Z}^+}$ be an $(\mathcal{F}_t)$ -adapted process on $[0, \infty)$. Suppose that for some $B \in (0, \infty)$, (4) holds for all $t \in \mathbb{Z}^+$. Then for any $\varepsilon > 0$, a.s., for all but finitely many $t \in \mathbb{Z}^+$,*

$$\max_{0 \leq s \leq t} X_s \leq t(\log t)^{1+\varepsilon}.$$

The next result is contained in Theorem 3.3 of [17].

Lemma 3 *Let $(X_t)_{t \in \mathbb{Z}^+}$ be an $(\mathcal{F}_t)$ -adapted process on $[0, \infty)$, such that for some $b \in \mathbb{N}$*

$$\mathbb{P}(X_{t+1} \leq X_t + b) = 1,$$

for all $t \in \mathbb{Z}^+$. Suppose that there exists $f : [0, \infty) \rightarrow [0, \infty)$ a nondecreasing function and $\delta > 0$ for which

$$\mathbb{E}[f(X_{t+1}) - f(X_t) \mid \mathcal{F}_t] \geq \delta \text{ a.s.}$$

for all $t \in \mathbb{Z}^+$. Suppose $\varepsilon > 0$. For $x \geq 0$ define the function r_ε by

$$r_\varepsilon(x) := \inf\{y \geq 0 : \delta^{-1}(\log y)^{1+\varepsilon} f(y+b) \geq x\}.$$

Then, a.s., for all but finitely many $t \in \mathbb{Z}^+$,

$$\max_{0 \leq s \leq t} X_s \geq r_\varepsilon(t) - b.$$

We also state the following result on existence of passage-time moments for one-dimensional stochastic processes, which is a simple consequence of Theorem 1 of [4].

Lemma 4 *Let $(X_t)_{t \in \mathbb{Z}^+}$ be an $(\mathcal{F}_t)_{t \in \mathbb{Z}^+}$ -adapted stochastic process taking values in an unbounded subset $\mathcal{S}$ of $[0, \infty)$. Suppose $B > 0$. Set*

$$v_B := \min\{t \in \mathbb{N} : X_t \leq B\}.$$

Suppose that there exist $C \in (0, \infty)$, $\gamma \in [0, 1)$ such that for any $t \in \mathbb{Z}^+$

$$\mathbb{E}[X_{t+1} - X_t \mid \mathcal{F}_t] \leq -CX_t^\gamma \quad \text{on } \{v_B > t\}.$$

Then for any $p \in [0, 1/(1-\gamma)]$, for any $x \in \mathcal{S}$, $\mathbb{E}[v_B^p \mid X_0 = x] < \infty$.

4.2 Exclusion-voter configurations

We introduce some more notation. For $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ and $i \in \{1, \dots, N\}$ let

$$R_i := R_i(S) := \sum_{j=1}^i n_j, \quad T_i := T_i(S) := \sum_{j=i}^N m_j. \quad (6)$$

It is convenient to represent a configuration $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ diagrammatically as a right-down path in the quarter-lattice $\mathbb{Z}^+ \times \mathbb{Z}^+$. This is done as follows. Starting from $(0, T_1)$, construct a walk by reading left-to-right the configuration S and for each 0 (1) taking a unit




Figure 2: An example staircase configuration.

step in the right (down) direction. Thus the walk always starts with a step to the right, and will end at $(R_N, 0)$ after $|S|$ steps in total. See the Figure 2 for the case of S as given by (1).

The lattice squares of $\mathbb{Z}^+ \times \mathbb{Z}^+$ bounded by the right-down path determined by S constitute a polygonal region in the plane that we call the *staircase* corresponding to S .

With this representation of the configuration-space, the exclusion-voter model can be viewed as a growth/depletion process on staircases. For instance, exclusion moves are particularly simple in this context, corresponding to adding or removing a square at a corner.

As well as the ground-state configuration $\mathcal{D}_0$, we will introduce special notation for one more configuration. Set

$$\mathcal{D}_1 := \dots 11101000 \dots, \quad (7)$$

the configuration with $N(\mathcal{D}_1) = 1$ and vector representation $(1, 1)$.

We now introduce notation for the changes in configuration brought about by voter and exclusion moves. Given the staircase of S , there are $2N + 1$ ‘corners’ representing 10s and 01s alternately, of which $N + 1$ are 10s and N are 01s. In the staircase representation, these corners have coordinates (R_i, T_{i+1}) , $i \in \{0, \dots, N\}$ (for 10s) and (R_i, T_i) , $i \in \{1, \dots, N\}$ (for 01s), where $R_0 = T_{N+1} = 0$. Enumerate the 10s left-to-right in the configuration S by $0, 1, \dots, N$, and similarly the 01s by $1, \dots, N$.

For $j \in \{0, 1, \dots, N\}$, let $v_j^{10 \rightarrow 00}(S)$, $v_j^{10 \rightarrow 11}(S)$ denote the configuration obtained from S by performing a voter move changing the j th 10 to 00, 11 respectively. Similarly for $j \in \{1, \dots, N\}$ let $v_j^{01 \rightarrow 00}(S)$, $v_j^{01 \rightarrow 11}(S)$ denote the configuration obtained from the two possible voter moves at the j th 01. We use analogous notation for exclusion moves: $e_j^{10 \rightarrow 01}(S)$ ($j \in \{0, \dots, N\}$), $e_j^{01 \rightarrow 10}(S)$ ($j \in \{1, \dots, N\}$).

To conclude this section, we prove Proposition 1. We do not give all the details (which are elementary).

Proof of Proposition 1. It is not hard to see that $\mathcal{D}_0$ is an absorbing state for the pure voter model ($\beta = 1$) and for the left-moving totally asymmetric exclusion process ($\beta = 0, p = 1$), and hence also for the mixture model under $\mathbb{P}_{\beta,1}$ for any $\beta \in [0, 1]$.

We now turn to the communication of states. To show that all states within $\mathcal{D}$ communicate, it suffices to show that $\mathbb{P}_{\beta,p}(\xi_{t+1} = S_1 \mid \xi_t = S_0) > 0$ for each of the following:

- (i) $S_0 = \mathcal{D}_0, S_1 = \mathcal{D}_1$;
- (ii) $S_0 = \mathcal{D}_1, S_1 = \mathcal{D}_0$;
- (iii) any S_0 with $|S_0| \geq 2$ and some S_1 with $|S_1| = |S_0| + 1$;
- (iv) any S_0 with $|S_0| \geq 3$ and some S_1 with $|S_1| = |S_0| - 1$;
- (v) any S_0 with $|S_0| \geq 3$ and any S_1 where S_1 is identical to S_0 apart from in a single position $j \in \{2, 3, \dots, |S_0| - 1\}$.

In other words, given that moves of types (i)–(v) can occur, it is possible to step between any two configurations in $\mathcal{D}$ by first adjusting the length of the configuration via moves of types (i)–(iv) and then flipping the states in the interior of the configuration via moves of type (v). Similarly, to show that all states in $\mathcal{D} \setminus \{\mathcal{D}_0\}$ communicate, it suffices to show that all moves of types (iii)–(v) have positive probability.

It is not hard to see that voter moves can perform moves of types (ii), (iii) and (iv). Similarly exclusion moves with $p < 1$ can perform moves of types (i) and (iii), while exclusion moves with $p > 0$ can perform moves of types (ii) and (iv).

We claim that moves of type (v) can be performed provided: (a) $\beta \in (0, 1)$; or (b) $\beta = 0$ and $p \in (0, 1)$.

In case (a), suppose we need to replace a 0 by a 1 in the interior of a given configuration. If $p < 1$, we may perform a voter move on the first 10 to the left of the position to be changed, and then, if necessary, perform successive $10 \mapsto 01$ exclusion moves to ‘step’ the 1 into the desired position. If $p > 0$, an analogous procedure works, starting from the first 01 to the *right*. On the other hand, if we need to replace a 1 by a 0, a similar argument applies.

In case (b) we cannot use voter moves, but both types of exclusion move are permitted, so we can ‘bring in’ any 0 or 1 from outside the disordered region and then rearrange as necessary.

It follows that moves of type (ii)–(v) are possible provided $\beta \neq 1$ and $(\beta, p) \notin \{(0, 0), (0, 1)\}$, and all (i)–(v) are possible if we additionally impose $p < 1$.

To complete the proof we need to demonstrate aperiodicity in the case where $\beta \neq 1$, $p < 1$ and $(\beta, p) \neq (0, 0)$, where all states communicate. Since $\beta \neq 1$, exclusion moves may occur. Moreover, every configuration other than $\mathcal{D}_0$ contains at least one pair of each type (01 and 10). Hence there is a positive probability that a configuration other than $\mathcal{D}_0$ remains unchanged at a given step (when a proposed exclusion move fails to occur). Thus, since all states communicate, we have aperiodicity. $\square$

4.3 Lyapunov function lemmas

Throughout this paper, Lyapunov-type functions for ξ will be primary tools. In this section we introduce some of our functions and give some preliminary results.

Recall the definitions of R_i, T_i from (6). In [5], considerable use was made of the functions f_1, f_2 defined as follows: for $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ set

$$f_1(S) := \sum_{i=1}^N m_i R_i = \sum_{i=1}^N n_i T_i; \quad f_2(S) := \frac{1}{2} \left(\sum_{i=1}^N m_i R_i^2 + \sum_{i=1}^N n_i T_i^2 \right);$$

and put $f_1(\mathcal{D}_0) = f_2(\mathcal{D}_0) = 0$. Note that with the diagrammatical representation described in Section 4.2, f_1 is the area of the staircase; e.g. for S given by (1), $f_1(S) = 81$.

In the present paper we introduce some more Lyapunov-type functions that will prove valuable: these include ρ^2 (see (46) below), ϕ_α for $\alpha > 0$ (see (22) below), and g which we define shortly. First we state a result on f_1, f_2 contained in Lemma 4.1 of [5].

Lemma 5 *For any $S \in \mathcal{D}$, we have*

$$\frac{1}{2}|S| \leq f_1(S) \leq \frac{1}{4}|S|^2; \quad (8)$$

$$\frac{1}{4}|S|^2 \leq f_2(S) \leq \frac{1}{8}|S|^3. \quad (9)$$

The next result compares f_1 with f_2 .

Lemma 6 *For any $S \in \mathcal{D}$ we have*

$$f_2(S) \leq |S|f_1(S) \leq 2(f_1(S))^2. \quad (10)$$

Proof. We have that for $S \in \mathcal{D}$

$$f_2(S) \leq \frac{1}{2} \left(\sum_{i=1}^N m_i R_i + \sum_{i=1}^N n_i T_i \right) \cdot (R_N + T_1) = f_1(S) \cdot |S|, \quad (11)$$

since, by (6), $R_i \leq R_N$ and $T_i \leq T_1$ for $1 \leq i \leq N$. Then from (11) and the first inequality in (8) we obtain (10). $\square$

Now we collect some formulae for the expected increments of $f_1(\xi_t)$ and $f_2(\xi_t)$, derived in [5], that we will need.

Lemma 7 *Suppose $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ and $\beta, p \in [0, 1]$. Then*

$$\mathbb{E}_{\beta,p}[f_1(\xi_{t+1}) - f_1(\xi_t) \mid \xi_t = S] = (1 - \beta) \frac{N(1 - 2p) + (1 - p)}{2N + 1} - \beta \frac{N}{2N + 1}. \quad (12)$$

Also

$$\mathbb{E}_{\beta,p}[f_2(\xi_{t+1}) - f_2(\xi_t) \mid \xi_t = S] = (1 - \beta) \left(\frac{1}{2} + \frac{(1/2) - p}{2N + 1} - \frac{2p - 1}{2N + 1} \sum_{i=1}^N (R_i + T_i) \right). \quad (13)$$

Proof. The equations (12) and (13) follow from (7.2) and (5.3), (6.3) in [5]. $\square$

Next we define the function g , which captures most of f_1 , in a sense made precise in Lemma 8 below. For $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$, let $K = K(S)$ be the smallest member of $\{1, \dots, N\}$ for which $R_K T_K = \max_{1 \leq k \leq N} \{R_k T_k\}$. Then for $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ set

$$X(S) := R_K; \quad Y(S) := T_K; \quad (14)$$

and put $X(\mathcal{D}_0) = Y(\mathcal{D}_0) = 0$. Then for $S \in \mathcal{D}$ we define

$$g(S) = X(S)Y(S) = \max_{1 \leq k \leq N} \{R_k T_k\}, \quad (15)$$

where here $\max \emptyset = 0$. With the representation described in Section 4.2, g is the area of the largest rectangle that can be inscribed in the staircase.

Lemma 8 *For any $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$,*

$$f_1(S) \geq g(S) \geq \frac{f_1(S)}{1 + \log f_1(S)}. \quad (16)$$

Proof. We start with a geometrical argument that will yield the stated results via the staircase representation of configurations S . Define $f_a(x) := a/x$ for $a > 0$ and $x > 0$. For $a > 0$ and $b \geq 1$, let $R(a, b)$ denote the region defined by

$$R(a, b) := \{(x, y) \in \mathbb{R}^2 : 0 \leq x \leq b, 0 \leq y \leq (a/x)\mathbf{1}_{\{x \geq 1\}} + a\mathbf{1}_{\{x < 1\}}\}.$$

Then, with $|\cdot|$ denoting Lebesgue measure on $\mathbb{R}^2$,

$$|R(a, b)| = a + \int_1^b (a/x) dx = a + a \log b.$$

Let $h : [0, \infty) \rightarrow [0, c]$ be a nonincreasing bounded function such that $h(x) = c$ for $0 \leq x < 1$, $h(d) = 0$, and $h(x) \geq 1$ for $0 \leq x < d$, where $c \geq 1$ and $d \geq 1$. Denote

$$M := M(h) := \{(x, y) \in \mathbb{R}^2 : 0 \leq x \leq d, 0 \leq y \leq h(x)\}.$$

Let $a_0 := \sup\{a > 0 : \{f_a(x) : x > 0\} \cap M \neq \emptyset\}$, i.e. the greatest value of a for which a curve $f_a(x)$ intersects region M . Then let x_0 be such that $f_{a_0}(x_0) \in M$. Let $B(M)$ denote the rectangle with vertices $(0, 0)$, $(x_0, 0)$, $(0, f_{a_0}(x_0))$, and $(x_0, f_{a_0}(x_0))$; then $|B(M)| = x_0(a_0/x_0) = a_0$. Moreover, it is clear that $B(M) \subseteq M$ and $M \subseteq R(a_0, d)$. So we have that

$$|B(M)| \leq |M| \leq |R(a_0, d)| = a_0(1 + \log d). \quad (17)$$

So, using the fact that $f_{a_0}(d) = a_0/d \geq 1$, we obtain from (17) that

$$1 \leq \frac{|M|}{|B(M)|} \leq 1 + \log d \leq 1 + \log a_0 = 1 + \log |B(M)|. \quad (18)$$

Now we translate the above argument into a proof of the statements in the lemma. Fix a configuration $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ with block representation $(n_1, m_1, \dots, n_N, m_N)$. For $x \geq 0$ define

$$j_S(x) := \max\{j \in \mathbb{Z}^+, j \leq N : \sum_{i=1}^j n_i \leq x\},$$

$$h_S(x) := \sum_{i=j_S(x)+1}^N m_i,$$

where we interpret an empty sum as zero. Set $c_S = \sum_{i=1}^N m_i$ and $d_S = \sum_{i=1}^N n_i$. Then $h_S(x) = c_S$ when $0 \leq x < 1$, since $n_1 \geq 1$. Also, $h_S(x) = 0$ for $x \geq d_S$ and $h_S(x) \geq m_N \geq 1$ for $0 \leq x < d_S$. So h_S is a function of the form of h in the first paragraph of the present proof. In particular $|M(h_S)| = f_1(S)$ and $|B(M(h_S))| = g(S)$. Thus (18) implies (16). $\square$

5 Non-existence of passage-time moments

For $t \in \mathbb{Z}^+$, let $\mathcal{F}_t$ denote the σ -field generated by ξ_t . Recall the definitions of $X(S), Y(S)$ from (14). For convenience of notation, set $X_t := X(\xi_t)$ and $Y_t := Y(\xi_t)$. We consider the auxiliary $(\mathcal{F}_t)$ -adapted process $(\tilde{\xi}_t)_{t \in \mathbb{Z}^+}$ defined by $\tilde{\xi}_t := (X_t, Y_t) = (X(\xi_t), Y(\xi_t))$; then $\tilde{\xi}_t$ takes values in the quarter-lattice $\mathbb{Z}^+ \times \mathbb{Z}^+$, and $\xi_t = \mathcal{D}_0$ if and only if $\tilde{\xi}_t = (0, 0)$.

Let $\sigma_{x,y}$ be the time for ξ_t to hit the ground configuration $\mathcal{D}_0$ (equivalently, the time taken for ξ_t to hit the origin $(0, 0)$) given the $\mathcal{F}_0$ -event $\{X(\xi_0) = x, Y(\xi_0) = y\}$.

The crucial ingredient to the proof of non-existence of moments will be the following result, which plays a similar rôle to Lemma 6.2 in [5] in the case of the pure voter model, and the more general Lemmas 3.1 and 2 in [13] and [4] respectively: the latter two results apparently do not apply to our process.

Lemma 9 *Suppose $p \leq 1/2$ and $\beta \in [0, 1]$. There exist $\delta > 0$, $\gamma > 0$ such that for all $x, y \in \mathbb{Z}^+$*

$$\mathbb{P}_{\beta,p}(\sigma_{x,y} \geq \delta(x^2 + y^2)) \geq \gamma, \quad (19)$$

and for all $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$

$$\mathbb{P}_{\beta,p}\left(\tau \geq \delta \frac{f_1(S)}{1 + \log f_1(S)} \mid \xi_0 = S\right) \geq \gamma. \quad (20)$$

We note that (20) is close to Conjecture 7.2 in [5]. The proof of Lemma 9 will be carried out in stages. The model in this case is a mixture of the voter model with the transient exclusion process. The main difficulty in establishing Lemma 9 is that while the process $f_2(\xi_t)$ is a submartingale, it does not have uniformly bounded jumps: although the voter model is symmetric it can effect large changes very quickly.

The next result, a corollary to Lemma 1, gives control over the size of the disordered region in the mixture process of voter model with *symmetric or recurrent* exclusion ($p \geq 1/2$).

Lemma 10 Suppose $p \geq 1/2$ and $\beta \in [0, 1]$. Then for all $t \in \mathbb{N}$

$$\mathbb{P}_{\beta,p} \left(\max_{0 \leq s \leq t} |\xi_s| \leq 2\sqrt{10t^{1/2}} \right) \geq 0.95 - \frac{f_2(\xi_0)}{10t}.$$

Proof. For $p \geq 1/2$ and $\beta \in [0, 1]$, we have from (13) that $f_2(\xi_t)$ satisfies

$$\mathbb{E}_{\beta,p}[f_2(\xi_{t+1}) - f_2(\xi_t) \mid \xi_t = S] \leq \frac{1}{2},$$

for all $S \in \mathcal{D}$. Applying Lemma 1 to $f_2(\xi_t)$ with $r = 10t$ and $B = 1/2$, (5) implies that

$$\mathbb{P}_{\beta,p} \left(\max_{0 \leq s \leq t} f_2(\xi_s) \leq 10t \right) \geq 1 - \frac{(t/2) + f_2(\xi_0)}{10t} = 0.95 - \frac{f_2(\xi_0)}{10t}.$$

Then using the fact that $|S| \leq 2(f_2(S))^{1/2}$ for any $S \in \mathcal{D}$ (by (9)), we obtain the result. $\square$

Suppose that $\xi_0 = S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ with corresponding $\tilde{\xi}_0 = (x_0, y_0) \in \mathbb{Z}^+ \times \mathbb{Z}^+$, i.e., $X(S_0) = x_0, Y(S_0) = y_0$. In order to enable us to identify positions within a configuration $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$, enumerate the positions in the hybrid zone left to right as $1, 2, \dots, |S|$.

Now we return to the voter plus transient ($p \leq 1/2$) exclusion model, and define an auxiliary *coloured* process as follows. Set

$$H := \sum_{i=1}^{K(S_0)} (n_i + m_i),$$

recalling the definition of $K(S_0)$ from just above (14); then position H in S_0 is necessarily occupied by a 0 and position $H + 1$ by a 1. We *colour* the x_0 0s that occupy positions in $\{1, 2, \dots, H\}$ and the y_0 1s that occupy positions in $\{H + 1, \dots, |S_0|\}$. All other particles are uncoloured. Intuitively, coloured particles can be thought of as ‘high energy’. Next we will define the evolution of the colouring corresponding to the process $(\xi_t)_{t \in \mathbb{Z}^+}$. We emphasize that the colouring is associated with the *particles* (i.e., 1s and 0s) rather than the *sites*.

The colour dynamics is as follows. Exclusion moves do not alter any colour, so that particles retain their colour-state after an exclusion move. Voter moves affect colouring as follows: in a pair 01 or 10 suppose that the 1 is coloured while the 0 is not; a voter move to pair 00 produces two uncoloured particles while a move to pair 11 produces two coloured particles. On the other hand, if in an unlike pair the 0 is coloured and the 1 not, a voter move to 00 produces two coloured particles and to 11 produces two uncoloured particles.

We note the following facts about the dynamics:

- (a) Uncoloured 1s remain to the left of any coloured 1s, and uncoloured 0s remain to the right of coloured 0s.
- (b) A necessary condition for the process to be in the ground configuration $\mathcal{D}_0$ is that the set of coloured particles consists only of a (possibly empty) block of coloured 1s at the left boundary of the hybrid zone and a (possibly empty) block of coloured 0s at the right boundary.

With $\xi_0 = S_0 \in \mathcal{D}$, for $t \in \mathbb{N}$ let ξ_t^* denote the configuration ξ_t with the associated colouring as determined by $(\xi_0, \dots, \xi_t)$ according to the mechanism just described.

Let $\mathcal{F}_t^*$ denote the σ -field generated by ξ_t^* . Define the $\mathcal{F}_t^*$ -measurable random variables ℓ_t and r_t as follows. Let ℓ_t be the position (measured from the left end of the hybrid zone) of the leftmost coloured 1 in ξ_t^* and r_t be the position of the rightmost coloured 0 in ξ_t^* ; initially $r_0 + 1 = \ell_0$ by construction.

As the process evolves, coloured 1s may end up to the left of coloured 0s. We define an auxiliary process $(\zeta_t)_{t \in \mathbb{Z}^+}$ to keep track of such configurations. Informally, when $\ell_t < r_t$, ζ_t will be the portion of ξ_t between positions ℓ_t and r_t . More formally, we introduce a holding state $\mathcal{D}_0^*$ and set $\zeta_t = \mathcal{D}_0^*$ if $\ell_t \geq r_t$. If $\ell_t < r_t$, the configuration ξ_t^* induces a finite string of 0s and 1s obtained by extracting the segment of ξ_t^* between positions ℓ_t and r_t (inclusive); this string we call ζ_t . Then $(\zeta_t)_{t \in \mathbb{Z}^+}$ is an $(\mathcal{F}_t^*)$ -adapted process with $\zeta_0 = \mathcal{D}_0^*$.

Note that, when it is not in state $\mathcal{D}_0^*$, ζ_t contains only *coloured* particles when colours are transposed from ξ_t^* . Now the idea is that when $p \leq 1/2$, ζ_t behaves like the mixture of voter and $p \geq 1/2$ exclusion, except that the presence of uncoloured particles in ξ_t^* causes it to ‘slow down’; thus we aim for a version of Lemma 10 in this case. This is the next result.

Lemma 11 *Suppose $p \leq 1/2$ and $\beta \in [0, 1]$. Then for all $t \in \mathbb{N}$*

$$\mathbb{P}_{\beta,p} \left(\max_{0 \leq s \leq t} |\zeta_s| \leq 2\sqrt{10}t^{1/2} \right) \geq 0.95.$$

Proof. We compare the process $(\zeta_t)_{t \in \mathbb{Z}^+}$ to an independent copy $\xi' = (\xi'_t)_{t \in \mathbb{Z}^+}$ of the process ξ . We define $f_2^*(\zeta_t)$ analogously to $f_2(\xi_t)$, but counting only the (coloured) particles in region ζ_t , i.e. coloured 1s to the left of coloured 0s and coloured 0s to the right of coloured 1s. Suppose that initially we were to permit

$$\xi_0^* = \mathcal{D}'_0 := \dots 000111 \dots$$

where all 0s and 1s are coloured, so that $\zeta_0 = \mathcal{D}_0^*$. Then by a simple reflection argument, the process $(\zeta_t)_{t \in \mathbb{Z}^+}$ embedded in $(\xi_t^*)_{t \in \mathbb{Z}^+}$ started from $\xi_0^* = \mathcal{D}'_0$ has the same distribution under $\mathbb{P}_{\beta,p}$ as the process $(\xi_t)_{t \in \mathbb{Z}^+}$ under $\mathbb{P}_{\beta,1-p}$ with initial state $\mathcal{D}_0$. So in particular Lemma 10 holds with ζ_t instead of ξ_t given the initial configuration $\mathcal{D}'_0$; then using the fact that $f_2^*(\zeta_0) = 0$ we obtain the claimed result in this case.

Now, the presence of uncoloured 1s to the left of coloured 1s or uncoloured 0s to the right of coloured 0s restricts the growth of $|\zeta_t|$; hence the claimed result also holds for any permissible initial configuration for ξ_0^* other than $\mathcal{D}'_0$. (One can argue rigorously by stochastic domination at this point.) $\square$

Proof of Lemma 9. We first prove the statement (19). Let $\chi_t := \chi(\xi_t^*)$ denote the number of coloured particles in ξ_t^* . Then $(\chi_t)_{t \in \mathbb{Z}^+}$ is $(\mathcal{F}_t^*)$ -adapted and $\chi_0 = \chi(\xi_0^*) = x_0 + y_0$. Also, given $\chi_t = n$ for $n \in \mathbb{N}$ we have that $\chi_{t+1} = n$ unless a voter move is performed on a pair with exactly one particle coloured, in which case χ_{t+1} takes values $n - 1, n + 1$ with equal probability. Also if $\chi_t = 0$ then $\chi_{t+1} = 0$ as well. Thus χ_t is a nonnegative $(\mathcal{F}_t^*)$ -martingale with uniformly bounded jumps. It follows from Doob’s submartingale inequality applied to

the nonnegative submartingale $(\chi_t - (x_0 + y_0))^2$, using the fact that $\mathbb{E}[(\chi_t - (x_0 + y_0))^2] \leq t$ by the orthogonality of martingale increments, that for any $z > 0$

$$\mathbb{P}_{\beta,p} \left(\max_{0 \leq s \leq t} |\chi_s - (x_0 + y_0)|^2 \geq z \right) \leq t/z,$$

so that for all $\theta > 0$ and $t \in \mathbb{Z}^+$

$$\mathbb{P}_{\beta,p} \left(\max_{0 \leq s \leq t} |\chi_s - (x_0 + y_0)| \leq \theta t^{1/2} \right) \geq 1 - (1/\theta^2).$$

In particular,

$$\mathbb{P}_{\beta,p} \left(\min_{0 \leq s \leq t} \chi_s \geq (x_0 + y_0) - 10t^{1/2} \right) \geq 0.99.$$

Taking $t = \delta^2(x_0^2 + y_0^2)$ for some $\delta > 0$, combining the last display with Lemma 11, we have that with probability at least 0.94 the two events

$$\left\{ \min_{0 \leq s \leq t} \chi_s \geq (x_0 + y_0) - 10\delta(x_0^2 + y_0^2)^{1/2} \geq (1 - 10\delta)(x_0 + y_0) \right\}$$

and

$$\left\{ \max_{0 \leq s \leq t} |\zeta_s| \leq 2\sqrt{10}\delta(x_0^2 + y_0^2)^{1/2} \leq 2\sqrt{10}\delta(x_0 + y_0) \right\}$$

both occur (noting that $(x_0^2 + y_0^2)^{1/2} \leq (x_0 + y_0)$). Choose δ small, say $\delta = 0.01$. Then with good probability the total number χ_s of coloured particles up to time t remains greater than $0.9(x_0 + y_0)$ while the central overlap region ζ_s of coloured particles between the leftmost coloured 1 and the rightmost coloured 0 remains shorter than $0.1(x_0 + y_0)$. Hence there must remain at least one coloured 0 to the left of any coloured 1 or one coloured 1 to the right of any coloured 0. By observation (b) above, this excludes the possibility of $\xi_s = \mathcal{D}_0$ for any $s \leq t$, where $t = \delta^2(x_0^2 + y_0^2)$. Thus we obtain (19).

To derive (20), we use the fact that $x_0^2 + y_0^2 \geq 2x_0y_0 = 2g(\xi_0)$ and now use (16). This proves the lemma. $\square$

Remark. $(\chi_t)_{t \in \mathbb{Z}^+}$ is a martingale with bounded jumps on $\mathbb{Z}^+$; thus if τ^χ is the time at which all colours disappear (i.e. $\chi_t = 0$), we have $\mathbb{P}(\tau^\chi < \infty) = 1$ but $\mathbb{E}[\tau^\chi] = \infty$ (e.g. by Theorems 2.2.1 and 2.2.6 in [8]). However, since $\chi_t = 0$ neither implies nor is implied by $\xi_t = \mathcal{D}_0$, we cannot use this to conclude about either recurrence or non-ergodicity of ξ .

Now we are nearly ready to complete the proofs of Theorems 2 and 3. The proofs proceed in a similar way to the proof of Theorem 6.1 in [5].

Proof of Theorem 2. Take $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$. Suppose, for the purpose of deriving a contradiction, that $\mathbb{E}_{\beta,p}[\tau^{1+\varepsilon} \mid \xi_0 = S_0] < \infty$ for some $\varepsilon > 0$. We start with the simple bound, valid for any $t \in \mathbb{Z}^+$,

$$\tau^{1+\varepsilon} \geq (t + \tau)^{1+\varepsilon} \mathbf{1}_{\{\tau \geq t\}}. \tag{21}$$

Let $\xi' = (\xi'_t)_{t \in \mathbb{Z}^+}$ be an independent copy of ξ and τ' be the corresponding independent copy of τ . Then for any $t \in \mathbb{Z}^+$, conditioning on ξ_t , taking expectations in (21), and using the Markov property at time t , we obtain

$$\mathbb{E}_{\beta,p}[\tau^{1+\varepsilon} \mid \xi_0 = S_0] \geq \mathbb{E}_{\beta,p}[\mathbb{E}_{\beta,p}[(t + \tau')^{1+\varepsilon} \mid \xi'_0 = \xi_t] \mathbf{1}_{\{\tau \geq t\}} \mid \xi_0 = S_0].$$

Hence by (20), there exist $\delta > 0$, $\gamma > 0$ such that for any $t \in \mathbb{Z}^+$

$$\mathbb{E}_{\beta,p}[\tau^{1+\varepsilon} \mid \xi_0 = S_0] \geq \gamma \mathbb{E}_{\beta,p}[(t + \delta(f_1(\xi_t))^{1-(\varepsilon/2)})^{1+\varepsilon} \mathbf{1}_{\{\tau \geq t\}} \mid \xi_0 = S_0].$$

It follows that for some $\varepsilon' \in (0, \varepsilon)$ and some $C \in (0, \infty)$

$$\mathbb{E}_{\beta,p}[\tau^{1+\varepsilon} \mid \xi_0 = S_0] \geq C \mathbb{E}_{\beta,p}[(f_1(\xi_t))^{1+\varepsilon'} \mathbf{1}_{\{\tau \geq t\}} \mid \xi_0 = S_0] = C \mathbb{E}_{\beta,p}[(f_1(\xi_{t \wedge \tau}))^{1+\varepsilon'} \mid \xi_0 = S_0],$$

for any $t \in \mathbb{Z}^+$, using the fact that a.s. $f_1(\xi_\tau) = f_1(\mathcal{D}_0) = 0$. That is, given $\xi_0 = S_0$, $(f_1(\xi_{t \wedge \tau}))^{1+\varepsilon'}$ is uniformly bounded in L^1 .

Hence the assumption that $\mathbb{E}_{\beta,p}[\tau^{1+\varepsilon} \mid \xi_0 = S_0] < \infty$ implies that on $\xi_0 = S_0$ the process $f_1(\xi_{t \wedge \tau})$ is uniformly integrable, and trivially that $\tau < \infty$ a.s.; thus as $t \rightarrow \infty$, $\mathbb{E}_{\beta,p}[f_1(\xi_{t \wedge \tau}) \mid \xi_0 = S_0] \rightarrow \mathbb{E}_{\beta,p}[f_1(\xi_\tau) \mid \xi_0 = S_0] = f_1(\mathcal{D}_0) = 0$.

However, for $p \leq 1/2$ and $\beta \leq (1 - 2p)/(2 - 2p)$, it follows from (12) that any $t \in \mathbb{Z}^+$ and any $S \in \mathcal{D}$

$$\mathbb{E}_{\beta,p}[f_1(\xi_{t+1}) - f_1(\xi_t) \mid \xi_t = S] \geq 0.$$

Then by the submartingale property we have that for all $t \in \mathbb{Z}^+$, $\mathbb{E}_{\beta,p}[f_1(\xi_{t \wedge \tau}) \mid \xi_0 = S_0] \geq f_1(S_0) > 0$. Thus we have the desired contradiction. $\square$

Proof of Theorem 3. Suppose that $S_0 \in \mathcal{D} \setminus \{\mathcal{D}_0\}$, and, for a contradiction, that $\mathbb{E}_{\beta,p}[\tau^{2+\varepsilon} \mid \xi_0 = S_0] < \infty$ for some $\varepsilon > 0$. Then, for any $t \in \mathbb{Z}^+$, similarly to the proof of Theorem 2,

$$\mathbb{E}_{\beta,p}[\tau^{2+\varepsilon} \mid \xi_0 = S_0] \geq \mathbb{E}_{\beta,p}[\mathbb{E}_{\beta,p}[(t + \tau')^{2+\varepsilon} \mid \xi'_0 = \xi_t] \mathbf{1}_{\{\tau \geq t\}} \mid \xi_0 = S_0].$$

Hence for $p \leq 1/2$, using (20),

$$\begin{aligned} \mathbb{E}_{\beta,p}[\tau^{2+\varepsilon} \mid \xi_0 = S_0] &\geq \gamma \mathbb{E}_{\beta,p}[(t + \delta(f_1(\xi_t))^{1-(\varepsilon/3)})^{2+\varepsilon} \mathbf{1}_{\{\tau \geq t\}} \mid \xi_0 = S_0] \\ &\geq C \mathbb{E}_{\beta,p}[(f_1(\xi_{t \wedge \tau}))^{2+\varepsilon'} \mid \xi_0 = S_0] \geq C \mathbb{E}_{\beta,p}[(f_2(\xi_{t \wedge \tau}))^{1+\varepsilon''} \mid \xi_0 = S_0], \end{aligned}$$

using (10) for the last inequality. Hence the process $f_2(\xi_{t \wedge \tau})$ is uniformly integrable, and thus as $t \rightarrow \infty$, $\mathbb{E}_{\beta,p}[f_2(\xi_{t \wedge \tau}) \mid \xi_0 = S_0] \rightarrow \mathbb{E}_{\beta,p}[f_2(\xi_\tau) \mid \xi_0 = S_0] = f_2(\mathcal{D}_0) = 0$.

However, for $p \leq 1/2$ and $\beta \in [0, 1]$, for all $S \in \mathcal{D}$ and all $t \in \mathbb{Z}^+$ it follows from (13) that

$$\mathbb{E}_{\beta,p}[f_2(\xi_{t+1}) - f_2(\xi_t) \mid \xi_t = S] \geq 0.$$

Hence for all $t \in \mathbb{Z}^+$, $\mathbb{E}_{\beta,p}[f_2(\xi_{t \wedge \tau}) \mid \xi_0 = S_0] \geq f_2(S_0) > 0$, leading to a contradiction as before. $\square$

6 Recurrence

We consider a new Lyapunov-type function that generalizes f_1 . For $\alpha \geq 0$, set $\phi_\alpha(\mathcal{D}_0) := 0$ and for $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ set

$$\phi_\alpha(S) := \sum_{i=1}^N \sum_{j=R_{i-1}+1}^{R_i} \sum_{k=1}^{T_i} \frac{1}{(j+k)^\alpha} = \sum_{i=1}^N \sum_{j=T_{i+1}+1}^{T_i} \sum_{k=1}^{R_i} \frac{1}{(j+k)^\alpha}; \quad (22)$$

here and throughout this section we use the conventions $R_0 := 0, R_{N+1} := R_N, T_0 := T_1, T_{N+1} := 0$. In particular it follows from (22) that when $\alpha = 0$

$$\phi_0(S) = \sum_{i=1}^N n_i T_i = \sum_{i=1}^N m_i R_i = f_1(S).$$

The next lemma gives an expression for the expected increments of ϕ_α .

Lemma 12 *Let $\beta \in [0, 1]$ and $p \in [0, 1]$. Then for any $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ and any $t \in \mathbb{Z}^+$*

$$\begin{aligned} & \mathbb{E}_{\beta,p}[\phi_\alpha(\xi_{t+1}) - \phi_\alpha(\xi_t) \mid \xi_t = S] \\ &= \frac{\beta}{2N+1} \left\{ N \sum_{j=1}^N \left(\frac{1}{(T_j + R_j + 1)^\alpha} - \frac{1}{(T_j + R_{j-1} + 1)^\alpha} \right) \right. \\ & \quad \left. - \frac{1}{2} \sum_{j=2}^N \frac{1}{(T_j + R_j + 1)^\alpha} - \frac{N+1}{2} \frac{1}{(R_N + 1)^\alpha} \right\} \\ & \quad + \frac{1-\beta}{2N+1} \left\{ -p \sum_{j=1}^N \frac{1}{(T_j + R_j)^\alpha} + (1-p) \sum_{j=0}^N \frac{1}{(T_{j+1} + R_j + 2)^\alpha} \right\}. \end{aligned} \quad (23)$$

Proof. Recalling the notation of Section 4.2, write

$$\begin{aligned} D_j^{v,10}(S) &:= \phi_\alpha(v_j^{10 \mapsto 00}(S)) + \phi_\alpha(v_j^{10 \mapsto 11}(S)) - 2\phi_\alpha(S) \quad (j \in \{0, \dots, N\}); \\ D_j^{v,01}(S) &:= \phi_\alpha(v_j^{01 \mapsto 00}(S)) + \phi_\alpha(v_j^{01 \mapsto 11}(S)) - 2\phi_\alpha(S) \quad (j \in \{1, \dots, N\}); \\ D_j^{e,10}(S) &:= \phi_\alpha(e_j^{10 \mapsto 01}(S)) - \phi_\alpha(S) \quad (j \in \{0, \dots, N\}); \\ D_j^{e,01}(S) &:= \phi_\alpha(e_j^{01 \mapsto 10}(S)) - \phi_\alpha(S) \quad (j \in \{1, \dots, N\}). \end{aligned}$$

Summing over all possible moves we have that

$$\begin{aligned} & \mathbb{E}_{\beta,p}[\phi_\alpha(\xi_{t+1}) - \phi_\alpha(\xi_t) \mid \xi_t = S] \\ &= \frac{\beta}{2N+1} \left\{ \frac{1}{2} \sum_{j=1}^N D_j^{v,01}(S) + \frac{1}{2} \sum_{j=0}^N D_j^{v,10}(S) \right\} \\ & \quad + \frac{1-\beta}{2N+1} \left\{ -p \sum_{j=1}^N D_j^{e,01}(S) + (1-p) \sum_{j=0}^N D_j^{e,10}(S) \right\}. \end{aligned} \quad (24)$$

Now we calculate expressions for the terms in (24). The reader might find it helpful to refer to a picture such as Figure 2 in Section 4.2 here. We have that for $j \in \{1, \dots, N\}$

$$D_j^{v,01}(S) = \sum_{i=1}^N \left(\frac{1}{(T_i + R_i + 1)^\alpha} - \frac{1}{(T_i + R_{i-1} + 1)^\alpha} \right) - \frac{1}{(T_j + R_j + 1)^\alpha} - \frac{1}{(T_{j+1} + R_{j+1} + 1)^\alpha}.$$

Also for $j \in \{0, 1, \dots, N\}$

$$\begin{aligned} D_j^{v,10}(S) &= \sum_{i=1}^N \left(\frac{1}{(T_i + R_i + 1)^\alpha} - \mathbf{1}_{\{i \leq j\}} \frac{1}{(T_i + R_{i-1} + 1)^\alpha} - \mathbf{1}_{\{i > j\}} \frac{1}{(T_{i+1} + R_i + 1)^\alpha} \right) \\ &= \sum_{i=1}^N \left(\frac{1}{(T_i + R_i + 1)^\alpha} - \frac{1}{(T_i + R_{i-1} + 1)^\alpha} \right) + \frac{1}{(T_{j+1} + R_j + 1)^\alpha} - \frac{1}{(R_N + 1)^\alpha}. \end{aligned}$$

Now we deal with the exclusion moves, which are simpler. We get

$$D_j^{e,01}(S) = -\frac{1}{(T_j + R_j)^\alpha}; \quad D_j^{e,10}(S) = \frac{1}{(T_{j+1} + R_j + 2)^\alpha}.$$

Thus we have that

$$\begin{aligned} & \frac{1}{2} \sum_{j=1}^N D_j^{v,01}(S) + \frac{1}{2} \sum_{j=0}^N D_j^{v,10}(S) \\ &= \frac{2N+1}{2} \sum_{j=1}^N \left(\frac{1}{(T_j + R_j + 1)^\alpha} - \frac{1}{(T_j + R_{j-1} + 1)^\alpha} \right) - \frac{1}{2} \sum_{j=1}^N \frac{1}{(T_j + R_j + 1)^\alpha} \\ & \quad - \frac{1}{2} \sum_{j=1}^N \frac{1}{(T_{j+1} + R_{j+1} + 1)^\alpha} + \frac{1}{2} \sum_{j=0}^N \frac{1}{(T_{j+1} + R_j + 1)^\alpha} - \frac{N+1}{2} \frac{1}{(R_N + 1)^\alpha} \\ &= \frac{2N+1}{2} \sum_{j=1}^N \left(\frac{1}{(T_j + R_j + 1)^\alpha} - \frac{1}{(T_j + R_{j-1} + 1)^\alpha} \right) - \sum_{j=1}^N \frac{1}{(T_j + R_j + 1)^\alpha} \\ & \quad + \frac{1}{2} \frac{1}{(T_1 + R_1 + 1)^\alpha} + \frac{1}{2} \sum_{j=1}^N \frac{1}{(T_j + R_{j-1} + 1)^\alpha} - \frac{N+1}{2} \frac{1}{(R_N + 1)^\alpha} \\ &= N \sum_{j=1}^N \left(\frac{1}{(T_j + R_j + 1)^\alpha} - \frac{1}{(T_j + R_{j-1} + 1)^\alpha} \right) - \frac{1}{2} \sum_{j=2}^N \frac{1}{(T_j + R_j + 1)^\alpha} \\ & \quad - \frac{N+1}{2} \frac{1}{(R_N + 1)^\alpha}, \end{aligned}$$

where, as usual, an empty sum is understood to be 0.

Combining all the computations, from (24) we obtain (23). $\square$

In particular, since $\phi_0(S) = f_1(S)$, the case $\alpha = 0$ of (23) gives

$$\mathbb{E}_{\beta,p}[\phi_0(\xi_{t+1}) - \phi_0(\xi_t) \mid \xi_t = S] = \frac{-\beta N}{2N+1} + \frac{1-\beta}{2N+1}(N(1-2p) + (1-p)),$$

which agrees with (12). Here we will be interested in the properties of ϕ_1 . First we give a lower bound for ϕ_1 .

Lemma 13 *For any $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$,*

$$\phi_1(S) \geq \log(|S|/4).$$

Proof. Suppose $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$. From (22) we have that

$$\phi_1(S) \geq \sum_{i=1}^N \sum_{j=R_{i-1}+1}^{R_i} \frac{1}{1+j} = \sum_{j=1}^{R_N} \frac{1}{j+1}.$$

Here, by monotonicity,

$$\sum_{j=1}^{R_N} \frac{1}{1+j} \geq \int_1^{R_N} \frac{dx}{1+x} \geq \log(R_N/2).$$

Similarly, we obtain from (22) that $\phi_1(S) \geq \log(T_1/2)$. It follows that

$$\phi_1(S) \geq \log(\max\{R_N, T_1\}/2) \geq \log(|S|/4),$$

completing the proof. $\square$

The following lemma is the key to this section.

Lemma 14 *Suppose $\beta \geq 4/7$ and $p \in [0, 1]$. Then for any $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$,*

$$\mathbb{E}_{\beta,p}[\phi_1(\xi_{t+1}) - \phi_1(\xi_t) \mid \xi_t = S] \leq 0.$$

Proof. The main idea of the proof is to verify the essentially combinatorial fact that for any $N \in \mathbb{N}$, and any positive integers $n_1, m_1, \dots, n_N, m_N$, the right-hand side of equation (25) below is nonnegative provided $\beta \geq 4/7$.

For ease of notation during this proof, set $\Delta(S) := \mathbb{E}_{\beta,p}[\phi_1(\xi_{t+1}) - \phi_1(\xi_t) \mid \xi_t = S]$. It is clear from (23) that $\Delta(S)$ is non-increasing in p , and so it suffices to consider the case $p = 0$. Recall that $R_0 = 0$, $R_j = n_1 + \dots + n_j$ and $T_j = m_j + \dots + m_N$, $T_{N+1} = 0$. Then (23) gives

$$\begin{aligned} \Delta(S) &= \mathbb{E}_{\beta,0}[\phi_1(\xi_{t+1}) - \phi_1(\xi_t) \mid \xi_t = S] \\ &= \frac{\beta}{2N+1} \left\{ N \sum_{j=1}^N \left(\frac{1}{T_j + R_j + 1} - \frac{1}{T_j + R_{j-1} + 1} \right) - \frac{1}{2} \sum_{j=2}^N \frac{1}{T_j + R_j + 1} \right. \\ &\quad \left. - \frac{N+1}{2} \frac{1}{R_N + 1} \right\} + \frac{1-\beta}{2N+1} \left\{ \sum_{j=0}^N \frac{1}{T_{j+1} + R_j + 2} \right\}. \end{aligned} \tag{25}$$

We rewrite (25) by setting $\gamma := (1-\beta)/\beta \in [0, \infty)$ to obtain

$$\frac{2N+1}{\beta} \Delta(S) = N \sum_{j=1}^N \left(\frac{1}{T_j + R_j + 1} - \frac{1}{T_j + R_{j-1} + 1} \right)$$

$$-\frac{1}{2} \sum_{j=2}^N \frac{1}{T_j + R_j + 1} - \frac{1}{2} \frac{N+1}{R_N + 1} + \gamma \sum_{j=1}^{N+1} \frac{1}{T_j + R_{j-1} + 2}. \quad (26)$$

We need to show that the right-hand side of (26) is non-positive. Since this quantity is nondecreasing in γ , it suffices to consider the case $\gamma = 3/4$, corresponding to $\beta = 4/7$. Set

$$\begin{aligned} \tilde{\Delta}(S) := N \sum_{j=1}^N \left(\frac{1}{T_j + R_j + 1} - \frac{1}{T_j + R_{j-1} + 1} \right) - \sum_{j=2}^N \frac{1/2}{T_j + R_j + 1} - \frac{1}{2} \frac{N+1}{R_N + 1} \\ + \frac{3}{4} \sum_{j=1}^{N+1} \frac{1}{T_j + R_{j-1} + 1}, \end{aligned}$$

so that, from (26), $\Delta(S) \leq \frac{\beta}{2N+1} \tilde{\Delta}(S)$.

Write $A_N := 1 + m_1 + m_2 + \dots + m_N$, $D_0 := 0$ and, for $i \in \{1, \dots, N\}$, $D_i := (n_1 - m_1) + \dots + (n_i - m_i)$, so that $R_{j-1} + T_j + 1 = A_N + D_{j-1}$. Then we have that

$$\begin{aligned} \tilde{\Delta}(S) &= N \sum_{j=1}^N \left(\frac{1}{A_N + D_{j-1} + n_j} - \frac{1}{A_N + D_{j-1}} \right) - \sum_{j=2}^N \frac{1/2}{A_N + D_{j-1} + n_j} \\ &\quad - \frac{(N+1)/2}{A_N + D_N} + \frac{3}{4} \sum_{j=1}^{N+1} \frac{1}{A_N + D_{j-1}} \\ &= \frac{1/2}{A_N + n_1} + \sum_{j=1}^N \left(\frac{N-1/2}{A_N + D_{j-1} + n_j} - \frac{N-3/4}{A_N + D_{j-1}} \right) - \frac{N/2 - 1/4}{A_N + D_N}. \end{aligned} \quad (27)$$

It is now convenient to introduce the notation for $k \in \{1, \dots, N\}$

$$H_k(S) := \sum_{j=1}^k \left(\frac{N-1/2}{A_N + D_{j-1} + n_j} - \frac{N-3/4}{A_N + D_{j-1}} \right) - \frac{(N+k-1)/4}{A_N + D_k}.$$

We now claim that if $N \geq 2$, for any $k \in \{2, \dots, N\}$,

$$H_k(S) \leq H_{k-1}(S). \quad (28)$$

Then we have from (27) and (28) that for $N \geq 2$

$$\begin{aligned} \tilde{\Delta}(S) &= \frac{1/2}{A_N + n_1} + H_N(S) \leq \frac{1/2}{A_N + n_1} + H_1(S) \\ &= \frac{1/2}{A_N + n_1} + \frac{N-1/2}{A_N + n_1} - \frac{N-3/4}{A_N} - \frac{N/4}{A_N + D_1} \\ &= \frac{N}{A_N + n_1} - \frac{N-3/4}{A_N} - \frac{N/4}{A_N + n_1 - m_1} \\ &\leq \frac{N}{A_N + n_1} - \frac{N-3/4}{A_N + n_1} - \frac{N/4}{A_N + n_1} = \frac{(3-N)/4}{A_N + n_1}. \end{aligned}$$

Thus $\tilde{\Delta}(S) \leq 0$, and hence $\Delta(S) \leq 0$ also, for all $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$ with $N(S) \geq 3$.

Let us now verify the claim (28). We have that for $k \geq 2$

$$\begin{aligned} H_k(S) &= \sum_{j=1}^{k-1} \left(\frac{N-1/2}{A_N + D_{j-1} + n_j} - \frac{N-3/4}{A_N + D_{j-1}} \right) + \frac{N-(1/2)}{A_N + D_{k-1} + n_k} \\ &\quad - \frac{N-(3/4)}{A_N + D_{k-1}} - \frac{(N+k-1)/4}{A_N + D_k} \\ &= \sum_{j=1}^{k-1} \left(\frac{N-1/2}{A_N + D_{j-1} + n_j} - \frac{N-3/4}{A_N + D_{j-1}} \right) + \left[\frac{(3N-k-1)/4}{A_N + D_{k-1} + n_k} - \frac{N-(3/4)}{A_N + D_{k-1}} \right] \\ &\quad + \left[\frac{(N+k-1)/4}{A_N + D_{k-1} + n_k} - \frac{(N+k-1)/4}{A_N + D_k} \right], \end{aligned}$$

where we have split the term with denominator $A_N + D_{k-1} + n_k$ into two parts. Note that for all j we have $A_N + D_{j-1} + n_j \geq A_N + D_{j-1}$ and also $A_N + D_{j-1} + n_j = A_N + D_j + m_j \geq A_N + D_j$. Therefore, applying these inequalities separately to the two terms in square brackets in the last display, we obtain

$$H_k(S) \leq \sum_{j=1}^{k-1} \left(\frac{N-1/2}{A_N + D_{j-1} + n_j} - \frac{N-3/4}{A_N + D_{j-1}} \right) + \left[\frac{(-N-k+2)/4}{A_N + D_{k-1}} \right] = H_{k-1}(S).$$

This proves the claim (28).

To complete the proof of the lemma, we show that $\Delta(S) \leq 0$ for $N(S) \in \{1, 2\}$ also. For $N = 1$, we have from the $\beta = 4/7$ case of (26) that

$$\begin{aligned} \frac{21}{4}\Delta(S) &= \frac{1}{1+m_1+n_1} - \frac{1}{1+m_1} - \frac{1}{1+n_1} + \frac{3/4}{2+n_1} + \frac{3/4}{2+m_1} \\ &= -\frac{n_1 m_1 (n_1 - m_1)^2 + 13(n_1 + m_1) + 4(1 + n_1 m_1) + 14(n_1^2 + m_1^2) + 5(n_1^3 + m_1^3)}{4(1+m_1+n_1)(1+m_1)(1+n_1)(2+m_1)(2+n_1)} \leq 0. \end{aligned}$$

Finally, for $N = 2$, from (26) again,

$$\begin{aligned} \frac{35}{4}\Delta(S) &= \frac{2}{m_1 + m_2 + n_1 + 1} - \frac{2}{m_1 + m_2 + 1} + \frac{3/2}{m_2 + n_1 + n_2 + 1} - \frac{2}{m_2 + n_1 + 1} \\ &\quad - \frac{3/2}{n_1 + n_2 + 1} + \frac{3/4}{m_1 + m_2 + 2} + \frac{3/4}{m_2 + n_1 + 2} + \frac{3/4}{n_1 + n_2 + 2} \\ &= -\frac{Q}{R} \end{aligned}$$

where $R = 4(m_1 + m_2 + n_1 + 1)(m_1 + m_2 + 1)(m_2 + n_1 + n_2 + 1)(m_2 + n_1 + 1)(n_1 + n_2 + 1)(m_1 + m_2 + 2)(m_2 + n_1 + 2)(n_1 + n_2 + 2)$ and

$$Q = m_1 n_1^4 (2m_1^2 + 5n_1^2 - m_1 n_1) + n_2 m_2^4 (2n_2^2 + 5m_2^2 - n_2 m_2) + 244 \text{ positive terms}$$

as can be readily checked in **Maple**, for instance. Since $2x^2 + 5y^2 - xy$ is always nonnegative, we conclude that $\Delta(S) \leq 0$ in this last case also. $\square$

Proof of Theorem 4. Lemma 14 shows that for $\beta \geq 4/7$, $(\phi_1(\xi_t))_{t \in \mathbb{Z}^+}$ is a supermartingale on $\xi_t \in \mathcal{D} \setminus \{\mathcal{D}_0\}$. Since, by Lemma 13, $\phi_1(S) \rightarrow \infty$ as $|S| \rightarrow \infty$, we can use Theorem 2.2.1 of [8] to complete the proof of the theorem. $\square$

7 Existence of passage-time moments

Our main tool in this section will be Lemma 4 applied with the Lyapunov function f_2 . Our first result is a bound on the expected increments of f_2 .

Lemma 15 *Suppose $\beta \in [0, 1)$ and $p > 1/2$. Then there exists $C \in (0, \infty)$ such that for all but finitely many $S \in \mathcal{D}$*

$$\mathbb{E}_{\beta,p}[f_2(\xi_{t+1}) - f_2(\xi_t) \mid \xi_t = S] \leq -C(f_2(S))^{1/6}.$$

Proof. First, as noted just below (5.3) in [5], $R_N + T_1 = |S|$ and $R_i + T_i \geq N$, so that

$$\sum_{i=1}^N (R_i + T_i) \geq \max\{|S|, N^2\} \geq N|S|^{1/2}. \quad (29)$$

Then from (13) with (29) it follows that for $p > 1/2$

$$\mathbb{E}_{\beta,p}[f_2(\xi_{t+1}) - f_2(\xi_t) \mid \xi_t = S] \leq \frac{1-\beta}{2} - (1-\beta)(2p-1)\frac{N|S|^{1/2}}{2N+1}.$$

Hence we have

$$\mathbb{E}_{\beta,p}[f_2(\xi_{t+1}) - f_2(\xi_t) \mid \xi_t = S] \leq \frac{1-\beta}{2} - \frac{1}{3}(1-\beta)(2p-1)|S|^{1/2}.$$

Then the result follows from (9). $\square$

Proof of Theorem 6. Applying Lemma 4 with $X_t = f_2(\xi_t)$ and using Lemma 15 shows that the hitting time of a finite subset of $\mathcal{D}$ has finite $(6/5)$ -th moment. Since $\mathcal{D}_0$ is accessible from any state, it follows that τ also has finite $(6/5)$ -th moment. $\square$

Remark. The exponent $1/6$ in Lemma 15 may not be best possible. However, the example configuration $n_1 = n_2 = \dots = n_{N-1} = 1, n_N = N^2$ and $m_1 = N^2, m_2 = \dots = m_N = 1$ shows that one cannot increase the exponent to more than $1/4$. Hence the method used in this section seems unable to prove existence of moments greater than $4/3$, and in particular it does not seem possible to attain $(3/2) - \varepsilon$ in this way (see the remark just after the statement of Theorem 6).

To prove Theorem 5 we will again apply Lemma 4, but this time we will take $X_t = f_2(\xi_t)^M$ for arbitrary $M \in [1, \infty)$. To study the increments of this process we recall some facts about f_2 under exclusion moves; compare (5.1) and (5.2) in [5]. We have that

$$f_2(e_j^{10 \mapsto 01}(S)) = f_2(S) + 1 + R_j + T_{j+1}, \quad (j \in \{0, \dots, N\}); \quad (30)$$

$$f_2(e_j^{01 \mapsto 10}(S)) = f_2(S) + 1 - R_j - T_j, \quad (j \in \{1, \dots, N\}), \quad (31)$$

where $R_0 := 0$ and $T_{N+1} := 0$. Now we will prove the following lemma.

Lemma 16 Suppose $\beta = 0$ and $p > 1/2$. Let $M \in (0, \infty)$. Then there exists $C \in (0, \infty)$ such that for all but finitely many $S \in \mathcal{D}$

$$\mathbb{E}_{0,p}[f_2(\xi_{t+1})^M - f_2(\xi_t)^M \mid \xi_t = S] \leq -C f_2(S)^{M-(5/6)}.$$

Proof. For the duration of this proof write $\Delta(S) := \mathbb{E}_{0,p}[f_2(\xi_{t+1})^M - f_2(\xi_t)^M \mid \xi_t = S]$. By summing over all the possible exclusion moves and using (30) and (31) we obtain

$$\begin{aligned} \Delta(S) &= \frac{1-p}{2N+1} \sum_{j=0}^N [(f_2(S) + 1 + R_j + T_{j+1})^M - f_2(S)^M] \\ &\quad + \frac{p}{2N+1} \sum_{j=1}^N [(f_2(S) + 1 - R_j - T_j)^M - f_2(S)^M]. \end{aligned} \quad (32)$$

Note that the $M = 1$ case of (32) agrees with (13). Here we have

$$(f_2(S) + 1 + R_j + T_{j+1})^M - f_2(S)^M = f_2(S)^M \left[\left(1 + \frac{1 + R_j + T_{j+1}}{f_2(S)} \right)^M - 1 \right].$$

Noting that $1 + R_j + T_{j+1} \leq 1 + |S| = O(f_2(S)^{1/2})$ for any j , by (9), we can apply Taylor's theorem to obtain

$$(f_2(S) + 1 + R_j + T_{j+1})^M - f_2(S)^M = f_2(S)^M \left[M \frac{1 + R_j + T_{j+1}}{f_2(S)} + O(f_2(S)^{-1}) \right].$$

Proceeding similarly for the terms in the second sum in (32), we obtain

$$\Delta(S) = \frac{M}{2N+1} f_2(S)^{M-1} \left[N + (1-p) + (1-2p) \sum_{j=1}^N (R_j + T_j) \right] + O(f_2(S)^{M-1}), \quad (33)$$

where the implicit constant in $O(\cdot)$ does not depend on S . Then from (33) with (29) and (9) we obtain that for some $C_1, C_2 \in (0, \infty)$

$$\Delta(S) \leq C_1 f_2(S)^{M-1} - C_2 f_2(S)^{M-(5/6)}.$$

This yields the result. $\square$

Proof of Theorem 5. Take $X_t = f_2(\xi_t)^M$ for some $M \geq 1$. Then we have from Lemma 16

$$\mathbb{E}[X_{t+1} - X_t \mid \xi_t = S] \leq -C X_t^{1-\frac{5}{6M}}$$

for all but finitely many S . Thus we can apply Lemma 4 to obtain $\mathbb{E}[\tau^{6M/5}] < \infty$. Since $M \geq 1$ was arbitrary, the theorem follows. $\square$

8 Size of the hybrid zone

In this section we prove the almost-sure bounds on the rate of growth of $|\xi_t|$ that we stated in Section 2.

Lemma 17 *Suppose $\beta \in [0, 1]$ and $p \in [0, 1]$. Then for any $\varepsilon > 0$, $\mathbb{P}_{\beta,p}$ -a.s., for all but finitely many t ,*

$$\max_{0 \leq s \leq t} f_1(\xi_s) \leq t(\log t)^{1+\varepsilon}.$$

Proof. From (12) we have that for any $S \in \mathcal{D}$

$$\mathbb{E}_{\beta,p}[f_1(\xi_{t+1}) - f_1(\xi_t) \mid \xi_t = S] \leq \frac{N(S) + 1}{2N(S) + 1} \leq 1.$$

Then we can apply Lemma 2 with $X_t = f_1(\xi_t)$ to obtain the result. $\square$

Proof of Theorem 7. Lemma 17 with the simple inequality $f_1(\xi_t) \geq N(\xi_t)^2/2$ implies the $p < 1/2$ case of (2). By (13) we have that for $p \geq 1/2$ and all $S \in \mathcal{D}$

$$\mathbb{E}_{\beta,p}[f_2(\xi_{t+1}) - f_2(\xi_t) \mid \xi_t = S] \leq \frac{1 - \beta}{2}.$$

Hence Lemma 2 with $X_t = f_2(\xi_t)$ yields, for any $\varepsilon > 0$, $\mathbb{P}_{\beta,p}$ -a.s.,

$$\max_{0 \leq s \leq t} f_2(\xi_s) \leq t(\log t)^{1+\varepsilon}, \tag{34}$$

for all but finitely many t . Then (3) follows from (34) with (9), and the $p \geq 1/2$ case of (2) follows from (34) with the simple inequality $f_2(\xi_t) \geq (N(\xi_t))^3/3$ (obtained by replacing each m_i and n_i by 1 in the definition of f_2). $\square$

For the remainder of this section, we concentrate on the pure exclusion process, i.e. when $\beta = 0$. Again the Lyapunov function f_1 will be a primary tool here; the next result describes its behaviour in this case. We use the abbreviation $N_t := N(\xi_t)$.

Lemma 18 *Suppose $\beta = 0$ and $p \in [0, 1]$. Then $f_1(\xi_t)$ has jumps in $\{-1, 0, +1\}$ with transition probabilities*

$$\begin{aligned} \mathbb{P}_{0,p}(f_1(\xi_{t+1}) - f_1(\xi_t) = 1 \mid \mathcal{F}_t) &= (1 - p) \frac{N_t + 1}{2N_t + 1} \geq \frac{1-p}{2}, \\ \mathbb{P}_{0,p}(f_1(\xi_{t+1}) - f_1(\xi_t) = -1 \mid \mathcal{F}_t) &= p \frac{N_t}{2N_t + 1} \leq \frac{p}{2}, \\ \mathbb{P}_{0,p}(f_1(\xi_{t+1}) - f_1(\xi_t) = 0 \mid \mathcal{F}_t) &= \frac{N_t + p}{2N_t + 1}. \end{aligned} \tag{35}$$

Hence for all $t \in \mathbb{Z}^+$,

$$f_1(\xi_t) \leq f_1(\xi_0) + t. \tag{36}$$

Moreover, when $p < 1/2$, for any $c \in (0, (1/2) - p)$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many t ,

$$f_1(\xi_t) \geq ct. \tag{37}$$

Proof. (35) follows from equations (5.5) and (5.6) in [5]. Then (36) is immediate. From (35), we have that ξ_t stochastically dominates $\xi_0 + \sum_{s=1}^t W_s$ where $W_1, W_2, \dots$ are i.i.d. random variables taking values $+1, 0, -1$ with probabilities $q/2, 1/2, p/2$ respectively. Hence the SLLN and the fact that $\mathbb{E}[W_1] = (1/2) - p$ yields (37) for $p < 1/2$. $\square$

Corollary 4 *Suppose $\beta = 0$ and $p < 1/2$. Then there exists $c'(p) > 0$ such that for any $c \in (0, c'(p))$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many t ,*

$$|\xi_t| \geq ct^{1/2}. \quad (38)$$

Suppose $\beta = 0$. Then there exists $C \in (0, \infty)$ such that for any $p \in [0, 1]$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$

$$N(\xi_t) \leq Ct^{1/2}. \quad (39)$$

Proof. The bound (38) follows from (37) together with (8); (39) follows from (36) with the simple inequality $f_1(\xi_t) \geq (N(\xi_t))^2/2$. $\square$

The next lemma gives some properties of the process $(|\xi_t|)_{t \in \mathbb{Z}^+}$. Recall the definition of configuration $\mathcal{D}_1$ from (7).

Lemma 19 *Suppose $\beta = 0$ and $p \in [0, 1]$. For any $t \in \mathbb{Z}^+$, we have that*

$$\mathbb{P}_{0,p}(|\xi_{t+1}| = 2 \mid \xi_t = \mathcal{D}_0) = 1 - \mathbb{P}_{0,p}(|\xi_{t+1}| = 0 \mid \xi_t = \mathcal{D}_0) = 1 - p, \quad (40)$$

and

$$\begin{aligned} \mathbb{P}_{0,p}(|\xi_{t+1}| = 3 \mid \xi_t = \mathcal{D}_1) &= \frac{2(1-p)}{3}, & \mathbb{P}_{0,p}(|\xi_{t+1}| = 2 \mid \xi_t = \mathcal{D}_1) &= \frac{1+p}{3}, \\ \mathbb{P}_{0,p}(|\xi_{t+1}| = 0 \mid \xi_t = \mathcal{D}_1) &= \frac{p}{3}. \end{aligned} \quad (41)$$

Also for any $t \in \mathbb{Z}^+$, conditional on $\xi_t \in \mathcal{D} \setminus \{\mathcal{D}_0, \mathcal{D}_1\}$, $|\xi_{t+1}| - |\xi_t|$ takes values only in $\{-1, 0, +1\}$. Moreover for any $S \in \mathcal{D} \setminus \{\mathcal{D}_0, \mathcal{D}_1\}$

$$\mathbb{P}_{0,p}(|\xi_{t+1}| = |\xi_t| + 1 \mid \xi_t = S) = \frac{2(1-p)}{2N(S) + 1}, \quad (42)$$

and

$$\mathbb{P}_{0,p}(|\xi_{t+1}| = |\xi_t| - 1 \mid \xi_t = S) = \frac{p}{2N(S) + 1} (\mathbf{1}_{\{n_1(S)=1\}} + \mathbf{1}_{\{m_{N(S)}(S)=1\}}) \leq \frac{2p}{2N(S) + 1}. \quad (43)$$

Proof. The statements (40) and (41) are straightforward. Suppose that $\xi_t = S$ for some $S \in \mathcal{D} \setminus \{\mathcal{D}_0, \mathcal{D}_1\}$. Then $|S| \geq 2$ and exclusion moves cannot effect a change of magnitude more than 1. We have that $|\xi_{t+1}| = |S| + 1$ if and only if we select (with probability $2/(2N(S) + 1)$) one of the two extreme 10 places, and then (with probability $1 - p$) we flip the 10 to a 01. Similarly, $|\xi_t|$ can decrease by 1 if and only if there exists a configuration $\dots 11101 \dots$ at the left end or a configuration $\dots 01000 \dots$ at the right end, and then we select the 01 and flip to 10. The statement of the lemma follows. $\square$

Consider the transient case $p \leq 1/2$. Note that

$$|\xi_t| \geq 2N(\xi_t). \quad (44)$$

Lemma 20 Suppose $\beta = 0$ and $p = 1/2$. Then

$$\mathbb{E}_{0,1/2}[|\xi_{t+1}|^3 - |\xi_t|^3 \mid \mathcal{F}_t] \geq 4 \text{ a.s.} \quad (45)$$

Proof. First we have from (40) and (41) that

$$\mathbb{E}_{0,1/2}[|\xi_{t+1}|^3 - |\xi_t|^3 \mid \xi_t = \mathcal{D}_0] = 4, \quad \mathbb{E}_{0,1/2}[|\xi_{t+1}|^3 - |\xi_t|^3 \mid \xi_t = \mathcal{D}_1] = 5.$$

Thus it remains to consider the case where $\xi_t = S$ for $S \in \mathcal{D} \setminus \{\mathcal{D}_0, \mathcal{D}_1\}$. Here, from (42) and (43), writing $N_t = N(\xi_t)$, we have

$$\begin{aligned} \mathbb{E}_{0,1/2}[|\xi_{t+1}|^3 - |\xi_t|^3 \mid \xi_t = S] &\geq \frac{1}{2N_t + 1}((|\xi_t| + 1)^3 - |\xi_t|^3) + \frac{1}{2N_t + 1}((|\xi_t| - 1)^3 - |\xi_t|^3) \\ &= \frac{1}{2N_t + 1}(6|\xi_t| + 2) \geq \frac{12N_t}{2N_t + 1} \geq 4, \end{aligned}$$

since by (44) $|\xi_t| \geq 2N_t$, and $N_t \geq 1$. $\square$

Proof of Theorem 9. The upper bound in the theorem is implied by (3). For the lower bound, use Lemma 3 with $f(x) = x^3$ and $X_t = |\xi_t|$. Then using (45) and the fact that $|\xi_t|$ has uniformly bounded jumps (see Lemma 19) we obtain the desired result. $\square$

We now work towards the upper bound for $|\xi_t|$ for $p \in [0, 1]$ given in Theorem 8. For $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$, define the function

$$\rho^2(S) := \sum_{i=1}^N m_i^2 + \sum_{i=1}^N n_i^2, \quad (46)$$

and set $\rho^2(\mathcal{D}_0) := 0$. Here are some bounds for $\rho^2(S)$.

Lemma 21 For any $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$,

$$|S| \leq \frac{1}{2N(S)}|S|^2 \leq \rho^2(S) \leq |S|^2. \quad (47)$$

Proof. Suppose that $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$. For the upper bound, we have

$$\rho^2(S) \leq \sum_{i=1}^N (m_i + n_i)^2 \leq \left(\sum_{i=1}^N (m_i + n_i) \right)^2 = |S|^2.$$

For the lower bound, we have from Jensen's inequality that

$$\frac{1}{N} \sum_{i=1}^N m_i^2 \geq \left(\frac{1}{N} \sum_{i=1}^N m_i \right)^2,$$

and similarly for the n_i ; hence

$$\rho^2(S) \geq \frac{1}{N}(R_N^2 + T_1^2) \geq \frac{1}{2N}(R_N + T_1)^2 = \frac{1}{2N}|S|^2 \geq |S|,$$

since $|S| \geq 2N$, completing the proof. $\square$

Lemma 22 Suppose $\beta = 0$ and $p \in [0, 1]$. For $t \in \mathbb{Z}^+$ we have

$$\mathbb{E}_{0,p}[\rho^2(\xi_{t+1}) - \rho^2(\xi_t) \mid \mathcal{F}_t] \leq 2 \text{ a.s.}; \quad (48)$$

moreover, for any $\varepsilon > 0$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$

$$\max_{0 \leq s \leq t} \rho^2(\xi_s) \leq t(\log t)^{1+\varepsilon}. \quad (49)$$

Proof. We start by proving (48). First we note that for any $t \in \mathbb{Z}^+$

$$\mathbb{E}_{0,p}[\rho^2(\xi_{t+1}) - \rho^2(\xi_t) \mid \xi_t = \mathcal{D}_0] = 2(1-p) \leq 2.$$

We next need to verify (48) for any configuration $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$.

To this end, let $\Delta_{1,i}(S)$ denote the change in $\rho^2(S)$ when a $01 \mapsto 10$ exclusion move is performed on the i -th 01 pair in S ($i = 1, \dots, N$). Similarly let $\Delta_{2,i}(S)$ denote the change in $\rho^2(S)$ when a $10 \mapsto 01$ exclusion move is performed on the i -th 10 pair ($i = 1, \dots, N+1$). In symbols, that is we set

$$\Delta_{1,i}(S) := \rho^2(e_i^{01 \mapsto 10}(S)) - \rho^2(S); \quad \Delta_{2,i}(S) := \rho^2(e_i^{10 \mapsto 01}(S)) - \rho^2(S),$$

with the notation of Section 4.2. Then

$$\mathbb{E}_{0,p}[\rho^2(\xi_{t+1}) - \rho^2(\xi_t) \mid \xi_t = S] = \frac{1}{2N+1} \left(p \sum_{i=1}^N \Delta_{1,i}(S) + q \sum_{i=1}^{N+1} \Delta_{2,i}(S) \right). \quad (50)$$

We compute the two sums on the right-hand side of (50) separately. Consider first all N possible exclusion moves $01 \mapsto 10$. It is easy to see, by inspecting separately the cases when $m_i = 1$ or $n_i = 1$, that

$$\begin{aligned} \Delta_{1,i}(S) &= -(2m_i - 2)\mathbf{1}_{\{m_i > 1\}} - (2n_i - 2)\mathbf{1}_{\{n_i > 1\}} + 2m_{i+1}\mathbf{1}_{\{n_i=1\}} + 2n_{i-1}\mathbf{1}_{\{m_i=1\}} \\ &= 2[2 - m_i - n_i + m_{i+1}\mathbf{1}_{\{n_i=1\}} + n_{i-1}\mathbf{1}_{\{m_i=1\}}], \end{aligned}$$

with the additional convention $n_0 = m_{N+1} = -1/2$, to make this formula correct for $i = 1$ and $i = N$ as well. Hence

$$\begin{aligned} \frac{1}{2} \sum_{i=1}^N \Delta_{1,i}(S) &= 2N - \sum_{i=1}^N (m_i + n_i) + \sum_{i=1}^N (m_{i+1}\mathbf{1}_{\{n_i=1\}} + n_{i-1}\mathbf{1}_{\{m_i=1\}}) \\ &= 2N - \sum_{i=1}^N (m_i + n_i) + \sum_{i=2}^{N+1} m_i \mathbf{1}_{\{n_{i-1}=1\}} + \sum_{i=0}^{N-1} n_i \mathbf{1}_{\{m_{i+1}=1\}} \\ &= 2N - m_1 - n_N - \frac{\mathbf{1}_{\{n_N=1\}} + \mathbf{1}_{\{m_1=1\}}}{2} - \sum_{i=2}^N m_i \mathbf{1}_{\{n_{i-1} > 1\}} - \sum_{i=1}^{N-1} n_i \mathbf{1}_{\{m_{i+1} > 1\}}. \end{aligned} \quad (51)$$

Similarly, a $10 \mapsto 01$ exclusion move on the i -th 10 pair from the left ($i = 1, 2, \dots, N+1$) contributes

$$\Delta_{2,i}(S) = 2[2 - m_i - n_{i-1} + m_{i-1}\mathbf{1}_{\{n_{i-1}=1\}} + n_i\mathbf{1}_{\{m_i=1\}}],$$

with the conventions here $n_0 = m_{N+1} = 1/2$ and $n_{N+1} = m_0 = 0$, to make this formula correct for $i = 1$ and $i = N + 1$. Therefore,

$$\frac{1}{2} \sum_{i=1}^{N+1} \Delta_{2,i}(S) = 2(N+1) - \frac{1}{2} - \sum_{i=1}^N m_i \mathbf{1}_{\{n_i > 1\}} - \frac{1}{2} - \sum_{i=1}^{N-1} n_i \mathbf{1}_{\{m_i > 1\}}. \quad (52)$$

Combining (51) and (52) with (50), we conclude that

$$\begin{aligned} \mathbb{E}_{0,p}[\rho^2(\xi_{t+1}) - \rho^2(\xi_t) \mid \xi_t = S] &= \frac{2}{2N+1} (2N+q) - \frac{p}{2N+1} (\mathbf{1}_{\{m_1=1\}} + \mathbf{1}_{\{n_N=1\}}) \\ &\quad - \frac{2}{2N+1} \sum_{i=1}^N (m_i[p\mathbf{1}_{\{n_{i-1}>1\}} + q\mathbf{1}_{\{n_i>1\}}] + n_i[p\mathbf{1}_{\{m_{i+1}>1\}} + q\mathbf{1}_{\{m_i>1\}}]) \leq 2 \text{ a.s.}, \end{aligned}$$

where in this formula we set $n_0 = m_{N+1} = +\infty$. This completes the proof of (48).

Finally, (49) follows from (48) with Lemma 2, taking $X_t = \rho^2(\xi_t)$. $\square$

Remark. There can be no uniform upper bound like (48) for the voter model: consider for instance the configuration S with $N(S) = 2$, $m_1 = m_2 = 1$ and n_1, n_2 large.

Suppose $p \in [0, 1]$. Then from (49), (39) and the middle inequality in (47) we obtain that for any $\varepsilon > 0$, $\mathbb{P}_{0,p}$ -a.s., for all but finitely many $t \in \mathbb{Z}^+$

$$\max_{0 \leq s \leq t} |\xi_s| \leq 2 \max_{0 \leq s \leq t} (N(\xi_s) \cdot \rho^2(\xi_s))^{1/2} \leq t^{3/4} (\log t)^{(1/2)+\varepsilon}.$$

In order to prove the upper bound in Theorem 8, we will give an argument that improves this bound from $t^{3/4}$ to $t^{2/3}$ (ignoring logarithmic terms). First we need a lemma.

Lemma 23 *Let $N \in \mathbb{N}$. Suppose that $n_1, n_2, \dots, n_N \geq 0$ and that for some $A, B > 0$*

$$\sum_{i=1}^N n_i^2 \leq A \quad \text{and} \quad \sum_{i=1}^N i n_i \leq B.$$

Then

$$\sum_{i=1}^N n_i \leq (6AB)^{1/3}.$$

Proof. We have

$$\begin{aligned} \left(\sum_{i=1}^N n_i \right)^3 &= \sum_{i=1}^N n_i^3 + 3 \sum_{i=1}^N n_i^2 \sum_{j \neq i} n_j + 3 \sum_{i=1}^N n_i \sum_{1 \leq j, k < i; j \neq k} n_j n_k \\ &\leq 3 \sum_{i=1}^N n_i^2 \sum_{i=1}^N n_i + 3 \sum_{i=1}^N n_i \left(\sum_{j=1}^{i-1} n_j \right)^2. \end{aligned}$$

Using Jensen's inequality for the final term, this last expression is bounded by

$$3 \sum_{i=1}^N n_i^2 \sum_{i=1}^N i n_i + 3 \sum_{i=1}^N i n_i \sum_{j=1}^{i-1} n_j^2 \leq 6AB,$$

by hypothesis. $\square$

Proof of Theorem 8. Part (i) of the theorem is (39), and the lower bound in part (ii) of the theorem is (38). We derive the upper bound in part (ii). Since each block of 1s has at least one element, observe that $\sum_{i=1}^N n_i(N-i) \leq f_1(S)$, and also $\sum_{i=1}^N n_i^2 \leq \rho^2(S)$. Thus Lemma 23 implies that for any $S \in \mathcal{D} \setminus \{\mathcal{D}_0\}$

$$\sum_{i=1}^N n_i \leq (6f_1(S)\rho^2(S))^{1/3} \leq 2(f_1(S)\rho^2(S))^{1/3},$$

and the same argument applies for $\sum_{i=1}^N m_i$. Hence for any $S \in \mathcal{D}$

$$|S| \leq 4(f_1(S)\rho^2(S))^{1/3}. \quad (53)$$

Taking $S = \xi_t$, we have $f_1(\xi_t) \leq C_1 t$ for all t and some $C_1 \in (0, \infty)$ by (36). Also for any $\varepsilon > 0$, $\mathbb{P}_{0,p}$ -a.s. $\rho^2(\xi_t) \leq C_2 t(\log t)^{1+\varepsilon}$ for all t by (49), for some $C_2 \in (0, \infty)$. Using these bounds in (53) completes the proof. $\square$

References

- [1] E. Andjel, P.A. Ferrari, and A. Siqueira, Law of large numbers for the simple exclusion process, *Stochastic Processes Appl.* **113** (2004) 217–233.
- [2] S. Aspandiiarov and R. Iasnogorodski, Tails of passage-time for non-negative stochastic processes and an application to stochastic processes with boundary reflection in a wedge, *Stochastic Processes Appl.* **66** (1997) 115–145.
- [3] S. Aspandiiarov and R. Iasnogorodski, General criteria of integrability of functions of passage-times for nonnegative stochastic processes and their applications, *Theory Probab. Appl.* **43** (1999) 343–369; translated from *Teor. Veroyatnost. i Primenen.* **43** (1998) 509–539 (in Russian).
- [4] S. Aspandiiarov, R. Iasnogorodski, and M. Menshikov, Passage-time moments for non-negative stochastic processes and an application to reflected random walks in a quadrant, *Ann. Probab.* **24** (1996) 932–960.
- [5] V. Belitsky, P.A. Ferrari, M.V. Menshikov, and S.Yu. Popov, A mixture of the exclusion process and the voter model, *Bernoulli* **7** (2001) 119–144.
- [6] P. Clifford and A. Sudbury, A model for spatial conflict, *Biometrika* **60** (1973) 581–588.

- [7] J.T. Cox and R. Durrett, Hybrid zones and voter model interfaces, *Bernoulli* **1** (1995) 343–370.
- [8] G. Fayolle, V.A. Malyshev, and M.V. Menshikov, Topics in the Constructive Theory of Countable Markov Chains, Cambridge University Press, 1995.
- [9] P.A. Ferrari and L.P.R. Pimentel, Competition interfaces and second class particles, *Ann. Probab.* **33** (2005) 1235–1254.
- [10] F.G. Foster, On the stochastic matrices associated with certain queuing processes, *Ann. Math. Statist.* **24** (1953) 355–360.
- [11] R.A. Holley and T.M. Liggett, Ergodic theorems for weakly interacting infinite systems and the voter model, *Ann. Probab.* **3** (1975) 643–663.
- [12] P. Jung, The noisy voter-exclusion process, *Stochastic Processes Appl.* **115** (2005) 1979–2005.
- [13] J. Lamperti, Criteria for stochastic processes II: passage-time moments, *J. Math. Anal. Appl.* **7** (1963) 127–145.
- [14] T.M. Liggett, Interacting Particle Systems, Springer, 1985.
- [15] T.M. Liggett, Stochastic Interacting Systems: Contact, Voter and Exclusion Processes, Springer, 1999.
- [16] I.M. MacPhee, M.V. Menshikov, and A.R. Wade, Exit times from cones for non-homogeneous random walk with asymptotically zero drift. Preprint (2008) available at <http://arxiv.org/abs/0806.4561>.
- [17] M.V. Menshikov, M. Vachkovskaia, and A.R. Wade, Asymptotic behaviour of randomly reflecting billiards in unbounded tubular domains, *J. Statist. Phys.* **132** (2008) 1097–1133.
- [18] A. Perrut, Hydrodynamic limits for a two-species reaction-diffusion process, *Ann. Appl. Probab.* **10** (2000) 163–191.
- [19] H. Rost, Non-equilibrium behaviour of a many particle process: density profile and local equilibria, *Z. Wahrsch. Verw. Gebiete* **58** (1981) 41–53.
- [20] A. Sturm and J.M. Swart, Tightness of voter model interfaces, *Elect. Comm. in Probab.* **13** (2008) 165–174.
- [21] D. Williams, Probability With Martingales, Cambridge University Press, 1991.